



Full Length Article

Summation identities for the Kummer confluent hypergeometric function ${}_1F_1(a; b; z)$ Gradimir V. Milovanović^{a,b,*}, Arjun K. Rathie^c, Nevena M. Vasović^d^a Serbian Academy of Sciences and Arts, 11000, Beograd, Serbia^b University of Niš, Faculty of Sciences and Mathematics, 18000, Niš, Serbia^c Dept. of Mathematics, Vedant College of Engineering & Technology, Rajasthan Technical University, TULSI, Bundi, 323021, India^d University of Kragujevac, Faculty of Hotel Management and Tourism, 36210, Vrnjačka Banja, Serbia

ARTICLE INFO

ABSTRACT

Keywords:

Gegenbauer polynomial
 Kummer confluent hypergeometric function
 Legendre duplication formula
 Orthogonal polynomials
 Summation identities

The role which hypergeometric functions have in the numerical and symbolic calculation, especially in the fields of applied mathematics and mathematical physics motivated research in this paper. In this note, a general formula for the sum, with the Kummer confluent hypergeometric function ${}_1F_1(a; b; z)$ is derived and given in terms of the function ${}_2F_2(a; b; z)$. The idea for this investigation comes from the theory of generalized Gauss-Rys quadrature formulas developed recently by (Milovanović, 2018) and (Milovanović and Vasović, 2022). Several results are obtained as special cases of the main result.

1. Introduction

The generalized hypergeometric function ${}_pF_q$ is defined by

$${}_pF_q(a_1, \dots, a_p; b_1, \dots, b_q; z) = \sum_{\nu=0}^{\infty} \frac{(a_1)_\nu \cdots (a_p)_\nu}{(b_1)_\nu \cdots (b_q)_\nu} \frac{z^\nu}{\nu!},$$

where the Pochhammer symbol $(\lambda)_\nu$ is given by

$$(\lambda)_\nu = \lambda(\lambda+1)\cdots(\lambda+\nu-1) = \frac{\Gamma(\lambda+\nu)}{\Gamma(\lambda)},$$

and $\Gamma(\lambda)$ is familiar Euler's gamma function

$$\Gamma(\lambda) = \int_0^\infty t^{\lambda-1} e^{-t} dt \quad \text{for } \operatorname{Re}(\lambda) > 0.$$

In Wolfram's Mathematica, the function ${}_pF_q$ is implemented as HypergeometricPFQ and it is suitable for both symbolic and numerical calculation (Wolfram, 2003). For $p = q + 1$, it has a branch cut discontinuity in the complex z plane running from 1 to ∞ . When $p \leq q$ this series converges for each $z \in \mathbb{C}$. For some recent results on this subject, especially on transformations, summations and other applications see (Milovanović et al., 2018; Milovanović and Rathie, 2019).

In the case $p = q = 1$ the function ${}_1F_1(a; b; z)$ is known as the Kummer

confluent hypergeometric function and it has the following integral representation (Olver et al., 2010, p. 326, p. 326)

$${}_1F_1(a; b; z) = \frac{\Gamma(b)}{\Gamma(a)\Gamma(b-a)} \int_0^1 t^{a-1} (1-t)^{b-a-1} e^{zt} dt, \quad \operatorname{Re}(b) > \operatorname{Re}(a) > 0. \quad (1)$$

In this note we obtain a general summation identity for the Kummer confluent hypergeometric function ${}_1F_1(a; b; z)$. The idea for this investigation comes from the theory of generalized Gauss-Rys quadrature formulas developed recently in (Milovanović, 2018; Milovanović and Vasović, 2022). In the next section we prove this general summation identity and consider several special cases, giving a corollary as a simplest case of the general result. Some conclusions and possible applications of the obtained results are mentioned at the end of this short note.

2. The main results: summation identities

For a general sum of the form

$$\sum_{\nu=0}^k (-1)^\nu \binom{k}{\nu} (\nu+\lambda+s+\epsilon)_{k-s} \left(\nu + \frac{1}{2} + \epsilon \right) {}_{s-1}F_1 \left(\nu+s-\frac{1}{2}+\epsilon; \nu+\lambda+s+\epsilon; -x \right), \quad (2)$$

* Corresponding author. Serbian Academy of Sciences and Arts, 11000, Beograd, Serbia.

E-mail address: gvm@mi.sanu.ac.rs (G.V. Milovanović).

where ϵ is 0 or 1, $\lambda > -1/2$ and $s, k \in \mathbb{N}$, in this note we prove explicit expression in terms of the hypergeometric function ${}_2F_2$.

Theorem 1. Let $\lambda > -1/2$, $k, s \in \mathbf{N}$ and $1 \leq s \leq k$. Then we have

$$\sum_{\nu=0}^k (-1)^\nu \binom{k}{\nu} (\nu + \lambda + s + \epsilon)_{k-s} \left(\nu + \frac{1}{2} + \epsilon \right) {}_{s-1}F_1 \left(\nu + s - \frac{1}{2} + \epsilon; \nu + \lambda + s + \epsilon; -x \right)$$

$$= \frac{(-1)^{s-1} k! \left(\lambda + \frac{1}{2} \right)_k x^{k-s+1}}{(k-s+1)! (\lambda + k + \epsilon)_{k+1}} {}_2F_2 \left(k + \frac{1+\epsilon}{2}, k + \frac{\epsilon}{2} + 1; k-s+2, 2k+\epsilon+\lambda+1; -x \right),$$

where ϵ is equal to 0 or 1.

Proof. We start from the formula for Gegenbauer polynomials (Prudnikov et al., 1986, p. 529, Eq. (10))

$$\int_0^a z^{\alpha-1} (a^2 - z^2)^{\lambda-1/2} e^{-xz^2} C_{2k+\epsilon}^{\lambda} \left(\frac{z}{a} \right) dz = \frac{(-1)^k a^{\alpha+2\lambda-1}}{2(2k+\epsilon)!} (2\lambda)_{2k+\epsilon} \left(\frac{1+\epsilon-\alpha}{2} \right)_k$$

$$\times \frac{\Gamma(\lambda + \frac{1}{2}) \Gamma(\frac{\alpha+\epsilon}{2})}{\Gamma(\frac{1+\alpha+\epsilon}{2} + \lambda + k)} {}_2F_2 \left(\frac{\alpha}{2}, \frac{\alpha+1}{2}; \frac{1+\alpha-\epsilon}{2} - k, \frac{1+\alpha+\epsilon}{2} + \lambda + k; -a^2 x \right), \quad (3)$$

which holds for $\epsilon = 0$ or 1, $a > 0$, $\lambda > -1/2$ and $\operatorname{Re}(\alpha) > -\epsilon$.

Let $s \in \mathbb{N}$ and $1 \leq s \leq k$. First, we consider the right-hand side in (3) for $a = 1$. For these values of s we put

$$\alpha = \begin{cases} 2(s+\gamma)-1, & \epsilon = 0, \\ 2(s+\gamma), & \epsilon = 1, \end{cases} \quad (4)$$

where $0 < |\gamma| \ll 1$. Since

$$\alpha = s - \frac{1}{2}(1-\epsilon) + \gamma, \quad \frac{\alpha+1}{2} = s + \frac{1}{2}\epsilon + \gamma, \quad \frac{\alpha+\epsilon}{2} = s - \frac{1}{2} + \epsilon + \gamma,$$

$$\frac{1+\alpha-\epsilon}{2} = s + \gamma, \quad \frac{1+\alpha+\epsilon}{2} = s + \epsilon + \gamma$$

and

$$\left(\frac{1+\epsilon-\alpha}{2} \right)_k = (-1)^k \left(\frac{1+\alpha-\epsilon}{2} - k \right)_k = (-1)^k (s-k+\gamma)_k,$$

we split now the right-hand side of (3) into two parts, replacing there the previous values for α ,

$$R_k^{(1)}(\alpha, \lambda, \epsilon) = \frac{(-1)^k (2\lambda)_{2k+\epsilon}}{2(2k+\epsilon)!} \frac{\Gamma(\lambda + \frac{1}{2}) \Gamma(\frac{\alpha+\epsilon}{2})}{\Gamma(\frac{1+\alpha+\epsilon}{2} + \lambda + k)}$$

$$= \frac{(-1)^k (2\lambda)_{2k+\epsilon}}{2(2k+\epsilon)!} \frac{\Gamma(\lambda + \frac{1}{2}) \Gamma(s - \frac{1}{2} + \epsilon + \gamma)}{\Gamma(s + \epsilon + \lambda + k + \gamma)}$$

and

$$R_k^{(2)}(\alpha, \lambda, \epsilon, x) = \left(\frac{1+\epsilon-\alpha}{2} \right)_k {}_2F_2 \left(\frac{\alpha}{2}, \frac{\alpha+1}{2}; \frac{1+\alpha-\epsilon}{2} - k, \frac{1+\alpha+\epsilon}{2} + \lambda + k; -x \right)$$

$$= (-1)^k (s-k+\gamma)_k$$

$$\times {}_2F_2 \left(s - \frac{1}{2}(1-\epsilon) + \gamma, s + \frac{1}{2}\epsilon + \gamma; s - k + \gamma, s + \epsilon + \lambda + k + \gamma; -x \right).$$

We note that

$$\hat{R}_k^{(1)}(2s-1+\epsilon, \lambda, \epsilon) = \lim_{\gamma \rightarrow 0} R_k^{(1)}(\alpha, \lambda, \epsilon) = \frac{(-1)^k (2\lambda)_{2k+\epsilon} \Gamma(\lambda + \frac{1}{2}) \Gamma(s - \frac{1}{2} + \epsilon)}{2(2k+\epsilon)!} \frac{\Gamma(s+\epsilon+\lambda+k)}{\Gamma(s+\epsilon+\lambda+k+k)}. \quad (5)$$

To find the corresponding value $\hat{R}_k^{(2)}(2s-1+\epsilon, \lambda, \epsilon, x) = \lim_{\gamma \rightarrow 0} R_k^{(2)}(\alpha, \lambda, \epsilon, x)$, we consider the series

$$(-1)^k (s-k+\gamma)_k {}_2F_2 \left(s - \frac{1}{2}(1-\epsilon) + \gamma, s + \frac{1}{2}\epsilon + \gamma; s - k + \gamma, s + \epsilon + \lambda + k + \gamma; -x \right)$$

$$= (-1)^k (s-k+\gamma)_k \sum_{\nu=0}^{\infty} \frac{\left(s - \frac{1}{2}(1-\epsilon) + \gamma \right)_\nu \left(s + \frac{1}{2}\epsilon + \gamma \right)_\nu}{(s-k+\gamma)_\nu (s+\epsilon+\lambda+k+\gamma)_\nu} \frac{(-x)^\nu}{\nu!}.$$

Because of

$$\lim_{\gamma \rightarrow 0} \frac{(s-k+\gamma)_k}{(s-k+\gamma)_\nu} = \begin{cases} 0, & \text{for } 0 \leq \nu \leq k-s, \\ \frac{(s-1)!}{(s-k+\nu-1)!}, & \text{for } k-s+1 \leq \nu \leq k, \end{cases}$$

we conclude that the first $k-s+1$ terms (for $\nu = 0, 1, \dots, k-s$) of this series, in the limit case, vanish, so that we consider the corresponding series with terms starting from the index $\nu = m = k-s+1$, i.e.,

$$(-1)^k (s-k+\gamma)_k \sum_{\nu=k-s+1}^{\infty} \frac{\left(s - \frac{1}{2}(1-\epsilon) + \gamma \right)_\nu \left(s + \frac{1}{2}\epsilon + \gamma \right)_\nu}{(s-k+\gamma)_\nu (s+\epsilon+\lambda+k+\gamma)_\nu} \frac{(-x)^\nu}{\nu!}$$

$$= (-1)^k (s-k+\gamma)_k \sum_{\nu=0}^{\infty} \frac{\left(s - \frac{1}{2}(1-\epsilon) + \gamma \right)_{\nu+m} \left(s + \frac{1}{2}\epsilon + \gamma \right)_{\nu+m}}{(s-k+\gamma)_{\nu+m} (s+\epsilon+\lambda+k+\gamma)_{\nu+m}} \frac{(-x)^{\nu+m}}{(\nu+m)!}.$$

Using the elementary identity $(p)_{\nu+m} = (p+m)_\nu (p)_m$ and letting $\gamma \rightarrow 0$, we get

$$\hat{R}_k^{(2)}(2s-1+\epsilon, \lambda, \epsilon, x) = (-1)^{s-1} \frac{(s-1)! \left(s - \frac{1}{2}(1-\epsilon) \right)_m \left(s + \frac{1}{2}\epsilon \right)_m}{(s+\epsilon+\lambda+k)_m}$$

$$x^m \sum_{\nu=0}^{\infty} \frac{\left(k + \frac{1+\epsilon}{2} \right)_\nu \left(k + 1 + \frac{\epsilon}{2} \right)_\nu}{(2k+\lambda+1+\epsilon)_\nu (\nu+m)!} \frac{(-x)^\nu}{\nu!}$$

$$= (-1)^{s-1} \frac{(s-1)! \left(s - \frac{1}{2}(1-\epsilon) \right)_{k-s+1} \left(s + \frac{1}{2}\epsilon \right)_{k-s+1}}{(s+\epsilon+\lambda+k)_{k-s+1} (k-s+1)!} x^{k-s+1}$$

$$\times {}_2F_2 \left(k + \frac{\epsilon}{2} + \frac{1}{2}, k + \frac{\epsilon}{2} + 1; k-s+2, 2k+\epsilon+\lambda+1; -x \right).$$

Now, we obtain the right-hand side of (3), under the assumed conditions, as a product of $R_k^{(1)}(2s-1+\epsilon, \lambda, \epsilon)$ and $\hat{R}_k^{(2)}(2s-1+\epsilon, \lambda, \epsilon, x)$:




Fig. 1. The sums (8) (left) and (9) (right) for $x \in [-1, 1]$, when $\lambda = 1/2$, $k = 4$ and $s = 1, 2, 3, 4$.

$$\begin{aligned} RHS &= \frac{(-1)^{k+s-1}(2\lambda)_{2k+\epsilon}}{2(2k+\epsilon)!} \frac{\Gamma(\lambda + \frac{1}{2})\Gamma(s - \frac{1}{2} + \epsilon)(s-1)!\left(s - \frac{1-\epsilon}{2}\right)_{k-s+1}\left(s + \frac{\epsilon}{2}\right)_{k-s+1}}{\Gamma(2k+\epsilon+\lambda+1)(k-s+1)!} \\ &\quad \times x^{k-s+1} {}_2F_2\left(k + \frac{\epsilon}{2} + \frac{1}{2}, k + \frac{\epsilon}{2} + 1; k - s + 2, 2k + \epsilon + \lambda + 1; -x\right). \end{aligned} \quad (6)$$

On the other side, for $a = 1$ and $z = \sqrt{t}$, the integral in (3) reduces to

$$\frac{1}{2} \int_0^1 t^{\alpha/2-1} (1-t)^{\lambda-1/2} e^{-xt} C_{2k+\epsilon}^\lambda(\sqrt{t}) dt,$$

which, after using the polynomial representation of the Gegenbauer polynomials

$$C_n^\lambda(x) = \sum_{\nu=0}^{\lfloor n/2 \rfloor} \frac{(-1)^\nu (\lambda)_{n-\nu}}{\nu!(n-2\nu)!} (2x)^{n-2\nu},$$

and taking $\alpha = 2s - 1 + \epsilon$ (as before in (4), with $\gamma = 0$), we get the following expression for the left-hand side of (3)

$$\begin{aligned} LHS &= \frac{1}{2} \sum_{\nu=0}^k \frac{(-1)^\nu (\lambda)_{2k-\nu+\epsilon} 2^{2(k-\nu)+\epsilon}}{\nu!(2k-2\nu+\epsilon)!} \int_0^1 t^{s-\frac{3}{2}+\epsilon+k-\nu} (1-t)^{\lambda-1/2} e^{-xt} dt \\ &= \frac{(-1)^k}{2} \sum_{\nu=0}^k \frac{(-1)^\nu (\lambda)_{k+\nu+\epsilon} 2^{2\nu+\epsilon}}{(k-\nu)!(2\nu+\epsilon)!} \int_0^1 t^{s-\frac{3}{2}+\epsilon+\nu} (1-t)^{\lambda-1/2} e^{-xt} dt. \end{aligned} \quad (7)$$

$$\begin{aligned} &\sum_{\nu=0}^k (-1)^\nu \binom{k}{\nu} (\nu + \lambda + s + \epsilon)_{k-s} \left(\nu + \frac{1}{2} + \epsilon\right)_{s-1} {}_1F_1\left(\nu + s - \frac{1}{2} + \epsilon; \nu + \lambda + s + \epsilon; -x\right) \\ &= \frac{(-1)^k 2k! \Gamma(\lambda)}{\sqrt{\pi} \Gamma(\lambda + \frac{1}{2})} \frac{(-1)^{k+s-1} (2\lambda)_{2k+\epsilon} \Gamma(\lambda + \frac{1}{2}) \Gamma(s - \frac{1}{2} + \epsilon) \Gamma(s) \left(s - \frac{1-\epsilon}{2}\right)_{k-s+1} \left(s + \frac{\epsilon}{2}\right)_{k-s+1}}{2\Gamma(2k+\epsilon+1) \Gamma(2k+\epsilon+\lambda+1) (k-s+1)!} \\ &\quad \times x^{k-s+1} {}_2F_2\left(k + \frac{\epsilon}{2} + \frac{1}{2}, k + \frac{\epsilon}{2} + 1; k - s + 2, 2k + \epsilon + \lambda + 1; -x\right) \\ &= \frac{(-1)^{s-1} k! \left(\lambda + \frac{1}{2}\right)_k x^{k-s+1}}{(k-s+1)! (\lambda + k + \epsilon)_{k+1}} {}_2F_2\left(k + \frac{\epsilon}{2} + \frac{1}{2}, k + \frac{\epsilon}{2} + 1; k - s + 2, 2k + \epsilon + \lambda + 1; -x\right), \end{aligned}$$

However, it can be expressed in terms of the Kummer confluent hypergeometric function (1) by taking $a = s - \frac{1}{2} + \epsilon + \nu$, $b = s + \nu + \lambda + \epsilon$, $z = -x$, so that

which had to be proved.

Theorem 1 for $\epsilon = 0$ and $\epsilon = 1$ gives the following sums:

Corollary 1. For $\lambda > -1/2$, $k, s \in \mathbf{N}$ and $1 \leq s \leq k$, we have

$$\sum_{\nu=0}^k (-1)^\nu \binom{k}{\nu} (\nu + \lambda + s)_{k-s} \binom{\nu + \frac{1}{2}}{s-1} {}_1F_1 \left(\nu + s - \frac{1}{2}; \nu + \lambda + s; -x \right) \\ = \frac{(-1)^{s-1} k! (\lambda + \frac{1}{2})_k x^{k-s+1}}{(k-s+1)! (\lambda+k)_{k+1}} {}_2F_2 \left(k + \frac{1}{2}, k+1; k-s+2, 2k+\lambda+1; -x \right) \quad (8)$$

and

$$\sum_{\nu=0}^k (-1)^\nu \binom{k}{\nu} (\nu + \lambda + s + 1)_{k-s} \binom{\nu + \frac{3}{2}}{s-1} {}_1F_1 \left(\nu + s + \frac{1}{2}; \nu + \lambda + s + 1; -x \right) \\ = \frac{(-1)^{s-1} k! (\lambda + \frac{1}{2})_k x^{k-s+1}}{(k-s+1)! (\lambda+k+1)_{k+1}} {}_2F_2 \left(k+1, k + \frac{3}{2}; k-s+2, 2k+\lambda+2; -x \right). \quad (9)$$

In the case $s = 1$ the function ${}_2F_2$ reduces to ${}_1F_1$, because $k-s+2 = k+1$, so that [Theorem 1](#) gives the following result:

Corollary 2. Let $\lambda > -1/2$ and $k \in \mathbb{N}$. Then we have

$$\sum_{\nu=0}^k (-1)^\nu \binom{k}{\nu} (\nu + \lambda + 1)_{k-1} {}_1F_1 \left(\nu + \frac{1}{2}; \nu + \lambda + 1; -x \right) \\ = \frac{(\lambda + \frac{1}{2})_k}{(k+\lambda)_{k+1}} x^k {}_1F_1 \left(k + \frac{1}{2}; 2k + \lambda + 1; -x \right)$$

and

$$\sum_{\nu=0}^k (-1)^\nu \binom{k}{\nu} (\nu + \lambda + 2)_{k-1} {}_1F_1 \left(\nu + \frac{3}{2}; \nu + \lambda + 2; -x \right) \\ = \frac{(\lambda + \frac{1}{2})_k}{(k+\lambda+1)_{k+1}} x^k {}_1F_1 \left(k + \frac{3}{2}; 2k + \lambda + 2; -x \right).$$

The sums from [Corollary 1](#) are displayed in [Fig. 1](#) as function in x , in the case $k = 4$ and $\lambda = 1/2$ and $s = 1, 2, 3, 4$.

Remark 1. The case $s > k$ is obvious. The general sum (2) should be written in the form

$$\sum_{\nu=0}^k (-1)^\nu \binom{k}{\nu} \frac{(\nu + \frac{1}{2} + \epsilon)_{s-1}}{(\nu + \lambda + k + \epsilon)_{s-k}} {}_1F_1 \left(\nu + s - \frac{1}{2} + \epsilon; \nu + \lambda + s + \epsilon; -x \right),$$

$\hat{R}_k^{(1)}(2s-1+\epsilon, \lambda, \epsilon)$ is given by (5), and $\hat{R}_k^{(2)}(2s-1+\epsilon, \lambda, \epsilon, x)$ is given directly by

$$(-1)^k (s-k)_k {}_2F_2 \left(s - \frac{1}{2}(1-\epsilon), s + \frac{1}{2}\epsilon; s-k, s+\epsilon+\lambda+k; -x \right).$$

3. Conclusion

In this paper, new summation identities for the Kummer confluent hypergeometric function have been obtained. As possible applications of given summation identities, we mention applications in representing results from the theory of orthogonal polynomials, theory of special functions, integral equations, etc. ([Mastroianni & Milovanović, 2008, 2009](#); [Asanov et al., 2017](#)).

Funding

The work of the first author was supported in part by the Serbian Academy of Sciences and Arts (Φ-96).

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Asanov, A., Matanova, K.B., Asanov, R.A., 2017. A class of linear and non linear Fredholm integral equations of the third kind. *Kuwait J. Sci.* 44 (1), 17–28.
- Mastroianni, G., Milovanović, G.V., 2008. Interpolation Processes – Basic Theory and Applications. Springer Monographs in Mathematics, Springer-Verlag, Berlin-Heidelberg.
- Mastroianni, G., Milovanović, G.V., 2009. Well-conditioned matrices for numerical treatment of Fredholm integral equations of the second kind. *Numer. Lin. Algebra Appl.* 16, 995–1011.
- Milovanović, G.V., 2018. An efficient computation of parameters in the Rys quadrature formula. *Bull. Cl. Sci. Math. Nat. Sci. Math.* 43, 39–64.
- Milovanović, G.V., Parmar, R.K., Rathie, A.K., 2018. A study of generalized summation theorems for the series ${}_2F_1$ with an applications to Laplace transforms of convolution type integrals involving Kummer's functions ${}_1F_1$. *Appl. Anal. Discrete Math.* 12, 257–272.
- Milovanović, G.V., Rathie, A.K., 2019. On a quadratic transformation due to Exton and its generalization. *Hacet. J. Math. Stat.* 48 (6), 1706–1711.
- Milovanović, G.V., Vasović, N., 2022. Orthogonal polynomials and generalized Gauss-Rys quadrature formulae. *Kuwait J. Sci.* 49 (1), 1–17.
- Olver, F.W.J., Lozier, D.W., Boisvert, R.F., Clark, C.W., 2010. NIST Handbook of Mathematical Functions. Cambridge University Press, Cambridge.
- Prudnikov, A.P., Brychkov, Y.A., Marichev, O.I., 1986. Special Functions. In: Integrals and Series, vol. 2. Gordon & Breach Science Publishers, New York.
- Wolfram, S., 2003. The Mathematica Book, the fifth ed. Wolfram Media, Inc.