

CALCULATING A CLASS OF INTEGRALS ENCOUNTERED IN
THEORETICAL CHEMISTRY

I. GUTMAN, G. V. MILOVANOVIĆ

(Presented at the 8th Meeting, held on October 30, 2009)

A b s t r a c t. A method for numerical calculating the Cauchy principal value integrals of the form $\text{v.p.} \int_{-\infty}^{+\infty} \log |P(ix)/Q(ix)| dx$ is developed, where $P(x)$ and $Q(x)$ are monic polynomials of equal degrees with integer coefficients, and $i = \sqrt{-1}$. These integrals play a distinguished role in theoretical chemistry.

AMS Mathematics Subject Classification (2000): 65D30, 92E10

Key Words: Numerical integration, theory of cyclic conjugation, cyclic conjugation energy effect, chemistry

1. Introduction

In this paper we are concerned with the Cauchy principal value integrals of the form

$$\text{v.p.} \int_{-\infty}^{+\infty} \log \left| \frac{P(ix)}{Q(ix)} \right| dx, \quad (1)$$

where

$$P(x) = \sum_{k=0}^n a_k x^k \quad \text{and} \quad Q(x) = \sum_{k=0}^n b_k x^k \quad (2)$$

are polynomials of equal degree whose coefficients are integers, $a_n = b_n = 1$, and $i = \sqrt{-1}$.

Integrals of this kind play a significant role in theoretical (quantum) chemistry. It seems that such integrals were first considered by Coulson and Jacobs [10], who showed that the difference between the total π -electron energy of two conjugated hydrocarbons with equal number of carbon atoms is given by

$$E(G_2) - E(G_1) = \frac{1}{\pi} \text{v.p.} \int_{-\infty}^{+\infty} \log \left| \frac{\phi(G_1, ix)}{\phi(G_2, ix)} \right| dx \quad (3)$$

where G_1 and G_2 are the corresponding molecular graphs [67], and ϕ stands for their characteristic polynomial. This formula is an immediate consequence of Coulson's classic integral expression for the total π -electron energy [9, 67]:

$$E(G) = \frac{1}{\pi} \text{v.p.} \int_{-\infty}^{+\infty} \left[n - \frac{ix \phi'(G, ix)}{\phi(G, ix)} \right] dx \quad (4)$$

where ϕ' denotes the first derivative of ϕ .

Several variants of Eq. (3), pertaining to energy differences, were considered in the chemical literature [18, 19, 28, 42]. Of these, the so-called "topological resonance energy" should especially be mentioned [60, 60]:

$$TRE(G) = \frac{1}{\pi} \text{v.p.} \int_{-\infty}^{+\infty} \log \left| \frac{\phi(G, ix)}{\alpha(G, ix)} \right| dx \quad (5)$$

where α is the matching polynomial.

It should be noted that in actual chemical applications (which are very numerous) both the total π -electron energy and the topological resonance energy are not computed by means of the formulas (3)–(5), but by using other computational techniques. However, there is another chemical theory in which calculation of numerical values of integrals of the type (1) cannot be avoided.

In 1977 one of the present authors [7, 17, 43] developed a novel theory of cyclic conjugation which made it possible to assess the effect of an individual

cycle on the thermodynamic stability of a polycyclic conjugated molecule. Details of this theory can be found in several expository articles [27, 36, 37, 38], whereas its mathematical formalism is outlined in [66, 67]. Almost in the same time Aihara [1] proposed a similar, yet not equivalent, theory, in which no integrals of the type (1) were used. The advantage of our approach over Aihara's was recognized only many years later [2]

Within our theory of cyclic conjugation, the energy-effect of a cycle Z of a polycyclic conjugated molecule whose molecular graph is G is computed as:

$$ef(G, Z) = \frac{1}{\pi} \text{v.p.} \int_{-\infty}^{+\infty} \log \left| \frac{\phi(G, ix)}{\phi(G, ix) + 2\phi(G - Z, ix)} \right| dx. \quad (6)$$

Recently, analogous expressions for the effects of pairs, triplets, quartets, etc. of cycles were deduced [74], as well as for the effect of conjugation in one cycle on conjugation in another cycle [91, 68, 16, 92, 39].

The quantity ef was extensively studied and applied to a variety of chemical problems. These researches were done either by finding some generally valid mathematical properties of ef [20, 21, 22, 24, 30, 65] or by performing numerical calculations [12, 25, 26, 29, 31, 32, 33, 34, 35, 41, 44, 45, 50, 51, 52, 53, 54, 56, 57, 61, 62, 63, 64, 69, 70, 71, 75, 72, 74, 76, 77, 78, 79, 5, 73, 81, 90, 4, 49, 48, 47, 3, 15, 55, 82, 46, 13, 14].

In the general case, the polynomials $P(x) \equiv \phi(G, x)$ and $Q(x) \equiv \phi(G, x) + 2\phi(G - Z, x)$, occurring in the expression on the right-hand side of Eq. (6) are monic, of equal degree, and have integer coefficients. The zeros of $Q(x)$ may be complex-valued and in practical applications are not known.

For $x \rightarrow \pm\infty$ the integrand in (6) tends to zero as $x^{-|Z|}$, where $|Z| \geq 3$ is the size of the cycle Z . At $x = 0$ the integrand may possess a singularity.

In standard chemical applications of the integrals of the form (1) it is assumed that the coefficients a_1 and b_1 in the polynomials $P(x)$ and $Q(x)$ are equal to zero. If this is not the case, then pertinent corrections need to be made [58, 85].

The hitherto reported ef -values were computed by means of a Simpson-type integration [40], in which the integrand is computed for $x = \frac{1}{2}h + kh$ for $k = 0, 1, 2, \dots$, up to the point at which the integrand is smaller than a critical value C . By empirical testing it was found that $h = 0.004$ and $C = 0.00001$ yield ef -values accurate to 3 or 4 decimal places. However, this latter accuracy could be tested only for the few (simple) examples for which the right-hand side of (6) can be solved analytically.

In the subsequent sections we show how integrals of the type (1) can be calculated in a much more efficient and much more accurate manner. Two methods are presented. In both cases a previous reduction to an integral with a rational function is provided (Section 2). In Section 3 we apply the trapezoidal rule after the so-called double exponential transformation of the integrand. Such ideas have been appeared in papers of Japanese mathematicians (cf. Takahasi and Mori [93, 94], Iri, Moriguti, and Takasawa [80], Mori [89]). The second method, presented in Section 4, is based on a transformation of the integral over the real line to an integral over the finite interval $(-1, 1)$, with respect to the Chebyshev weights. An application of the corresponding quadratures of Gaussian type is also presented.

2. Reduction to integrals of rational functions

In this section we reduce the Cauchy principal value integral (1) to an improper integral of a rational function over \mathbb{R} .

Let \mathcal{P}_n be a set of all real algebraic polynomials of degree at most n and $\hat{\mathcal{P}}_n$ be its subset of monic polynomials of degree n . With $\mathcal{R}[m, n]$ we denote the set of all rational functions of the form $u(t)/v(t)$ such that $u \in \mathcal{P}_m$, $v \in \hat{\mathcal{P}}_n$, and $\gcd(u(t), v(t)) = 1$ (i.e., the polynomials $u(t)$ and $v(t)$ are relatively prime).

According to (2) we have

$$P(ix)P(-ix) = (a_0 - a_2x^2 + a_4x^4 - \dots)^2 + x^2(a_1 - a_3x^2 + a_5x^4 - \dots)^2,$$

i.e., $|P(ix)|^2 = P(ix)P(-ix) = p(x^2)$ and similarly $|Q(ix)|^2 = Q(ix)Q(-ix) = q(x^2)$. Such polynomials

$$p(t) = t^n + \alpha t^{n-1} + \dots \quad \text{and} \quad q(t) = t^n + \beta t^{n-1} + \dots$$

are real monic polynomials of degree n and nonnegative for $t \geq 0$. Without loss of generality, we can suppose that they are relatively prime. In addition, we suppose that they have not positive zeros, i.e., $p(t), q(t) > 0$ for $t > 0$.

Then

$$\text{v.p.} \int_{-\infty}^{+\infty} \log \left| \frac{P(ix)}{Q(ix)} \right| dx = \frac{1}{2} \text{v.p.} \int_{-\infty}^{+\infty} \log \left| \frac{P(ix)}{Q(ix)} \right|^2 dx = \frac{1}{2} \text{v.p.} \int_{-\infty}^{+\infty} \log \frac{p(x^2)}{q(x^2)} dx.$$

An integration by parts gives

$$\frac{1}{2} \int \log \frac{p(x^2)}{q(x^2)} dx = \frac{x}{2} \log \frac{p(x^2)}{q(x^2)} - \int x^2 \left(\frac{p'(x^2)}{p(x^2)} - \frac{q'(x^2)}{q(x^2)} \right) dx. \quad (7)$$

Defining a rational function $R(t)$ by

$$R(t) = t \left(\frac{q'(t)}{q(t)} - \frac{p'(t)}{p(t)} \right) = \frac{h(t)}{p(t)q(t)}, \quad (8)$$

where $h(t) = t(p(t)q'(t) - q(t)p'(t))$, we get

$$\begin{aligned} R(t) &= t \left\{ \frac{nt^{n-1} + (n-1)\beta t^{n-2} + \dots}{t^n + \beta t^{n-1} + \dots} - \frac{nt^{n-1} + (n-1)\alpha t^{n-2} + \dots}{t^n + \alpha t^{n-1} + \dots} \right\} \\ &= t \frac{(\alpha - \beta)t^{2n-2} + \dots}{p(t)q(t)} \in \mathcal{R}[2n-1, 2n]. \end{aligned}$$

Precisely, $R(t) \in \mathcal{R}[m, 2n]$ for some m between 0 and $2n-1$.

Thus, $R(x^2) = o(x^{-2r})$ as $x \rightarrow +\infty$, where $r \geq 1$. For $t > 0$ the denominator of the rational function $R(t)$ is strictly positive, except $t = 0$, but this function $R(t)$ cannot have a pole at the origin. Namely, if $p(t)$ or $q(t)$ (not both) has a zero at the origin, then such a pole is eliminated by the factor t in $h(t)$. Regarding these facts, we have the existence of the improper integral

$$\int_{-\infty}^{+\infty} R(x^2) dx.$$

Also, by definition of the Cauchy principal value integral, it is easy to conclude that the first term on the right hand side in (7) has no contribution in the integral (1), so that the following result holds:

Lemma 1. *For the integral (1) we have*

$$\text{v.p.} \int_{-\infty}^{+\infty} \log \left| \frac{P(ix)}{Q(ix)} \right| dx = \int_{-\infty}^{+\infty} R(x^2) dx, \quad (9)$$

where the function $R(t) \in \mathcal{R}[m, 2n]$ ($0 \leq m \leq 2n-1$) has the form (8), where $h \in \mathcal{P}_m$ and $p, q \in \hat{\mathcal{P}}_n$.

For example, for $p(t) = t^2 + 4t$ and $q(t) = t^2 + 4t + 2$, according to (8) and (9), we get

$$\text{v.p.} \int_{-\infty}^{+\infty} \log \frac{x^4 + 4x^2}{x^4 + 4x^2 + 2} dx = \int_{-\infty}^{+\infty} \frac{-8(x^2 + 2)}{(x^2 + 4)(x^4 + 4x^2 + 2)} dx.$$

Thus, it is reduced to an improper integral, which value is

$$-2\pi \left(\sqrt{4 + 2\sqrt{2}} - 2 \right) = -3.852383833273321 \dots$$

Thus, our starting problem (1) is reduced to an integration of rational functions over \mathbb{R} . It is well known that in the eighteenth century Johan Bernoulli solved the problem of indefinite integration of rational functions by their partial decomposition. The main computational problem with this method is computing the factorization of a polynomial. However, in the middle of the nineteenth century the Russian mathematician Mikhail Vasilyevich Ostrogradsky presented an algorithm for finding the rational part of the integral without factoring. Some similar approaches were later discovered. The problem of computing the transcendental part of the primitive was recently solved. The recent development of symbolic computations made also a progress in this area (for details see a book of Bronstein [8], as well as some papers dealing with Landen transformation for rational functions [6] and [83]).

For our specific kind of integrals in the subsequent sections we give two efficient methods for their numerical calculating.

3. Double exponential transformation and trapezoidal rule

We start this section with some classical rules for calculating the integral $I(f) := \int_a^b f(x) dx$.

Taking $h := (b - a)/n$ and equally spaced points $x_k := a + kh$, $k = 0, 1, \dots, n$, we have the well-known composite trapezoidal rule

$$I(f) \approx T_n(f; h) := h \left(\frac{1}{2}f_0 + f_1 + \dots + f_{n-1} + \frac{1}{2}f_n \right), \quad (10)$$

where $f_k := f(x_k)$, $k = 0, 1, \dots, n$. If $f \in C^2[a, b]$ it is easy to prove that

$$I(f) - T_n(f; h) = -\frac{(b-a)h^2}{12} f''(\xi), \quad a < \xi < b. \quad (11)$$

As we can see this rule converges very slowly with respect to step refinement as $O(h^2)$.

Another simple rule is the classical composite Simpson rule

$$I(f) \approx S_n(f; h) := \frac{h}{3} [f_0 + 4(f_1 + \dots + f_{2n-1}) + 2(f_2 + \dots + f_{2n-2}) + f_{2n}],$$

where $h := (b - a)/2n$, $x_k := a + kh$, $f_k := f(x_k)$, $k = 0, 1, \dots, 2n$, which is slightly faster, but complicated than the previous one. Namely, if $f \in C^4[a, b]$,

$$I(f) - S_n(f; h) = -\frac{(b-a)h^4}{180} f^{(4)}(\xi), \quad a < \xi < b.$$

For functions with continuous derivatives of order at least $2m - 1$, a generalization of (11) is the well-known Euler-Maclaurin summation formula

$$\begin{aligned} I(f) - T_n(f; h) &= -\frac{h^2}{12}(f'(b) - f'(a)) + \frac{h^4}{720}(f'''(b) - f'''(a)) \\ &\quad - \dots - \frac{h^{2m} B_{2m}}{(2m)!} (f^{(2m-1)}(b) - f^{(2m-1)}(a)) - E_m(f), \end{aligned}$$

where B_{2m} is the Bernoulli number of order $2m$ and

$$E_m(f) = (b-a) \frac{B_{2m+2} h^{2m+2}}{(2m+2)!} f^{(2m+2)}(\xi), \quad a < \xi < b.$$

If we restrict our analysis to analytic functions with all derivatives of f which vanish at $x = a$ and $x = b$, then the discretization error is given only by remainder $E_m(f)$ as $m \rightarrow +\infty$. Then the convergence with respect to step refinement is faster than any finite order and the trapezoidal rule becomes a method of choice. Such a convergence is known as *exponential convergence*.

In order to calculate the integral (9) with the trapezoidal rule with the previous property we first apply the so-called double-exponential transformation $x = u(t) = \sinh((\pi/2) \sinh t)$, reducing it to

$$I = \int_{-\infty}^{+\infty} R(x^2) dx = \int_{-\infty}^{+\infty} R(u(t)^2) u'(t) dt,$$

i.e.,

$$I = \frac{\pi}{2} \int_{-\infty}^{+\infty} R\left(\sinh^2\left(\frac{\pi}{2} \sinh t\right)\right) \cosh\left(\frac{\pi}{2} \sinh t\right) \cosh t dt. \quad (12)$$

The crucial point in this transformation is the decay of the integrand be double exponential, i.e.,

$$|R(u(t)^2)u'(t)| \approx \exp(-C \exp |t|) \quad \text{as } |t| \rightarrow +\infty,$$

where C is some positive constant. For an integral of such form of an analytic function on $(-\infty, +\infty)$, it is known that the trapezoidal formula with an equal mesh size gives an optimal formula (cf. [80, 87, 88, 89, 93, 94, 95]).

In our case we apply the trapezoidal formula with an equal mesh size h , so that we obtain

$$I_h = \frac{\pi h}{2} \sum_{k=-\infty}^{+\infty} R\left(\sinh^2\left(\frac{\pi}{2} \sinh kh\right)\right) \cosh\left(\frac{\pi}{2} \sinh kh\right) \cosh kh.$$

Since the integrand decays double exponentially, in actual computation of the previous sum we truncate the infinite summation at $k = -M$ and $k = M$, so that we obtain the double-exponential (DE) formula for our integral

$$I \approx I_h^{(N)} = \frac{\pi h}{2} \sum_{k=-M}^M R\left(\sinh^2\left(\frac{\pi}{2} \sinh kh\right)\right) \cosh\left(\frac{\pi}{2} \sinh kh\right) \cosh kh,$$

where $N = 2M + 1$.

Example 1. Let $\phi(G, x) = x^{10} - 11x^8 + 41x^6 - 65x^4 + 43x^2 - 9$ and $\phi(G - Z, x) = x^4 - 3x^2 + 1$. Then we have $P(x) = \phi(G, x)$ and $Q(x) = \phi(G, x) + 2\phi(G - Z, x)$, so that


$$\begin{aligned} |P(ix)|^2 &= P(ix)P(-ix) = (9 + 43x^2 + 65x^4 + 41x^6 + 11x^8 + x^{10})^2 \\ |Q(ix)|^2 &= Q(ix)Q(-ix) = (7 + 37x^2 + 63x^4 + 41x^6 + 11x^8 + x^{10})^2, \end{aligned}$$

i.e.,

$$\left|\frac{P(ix)}{Q(ix)}\right|^2 = \left(\frac{9 + 16x^2 + 8x^4 + x^6}{7 + 16x^2 + 8x^4 + x^6}\right)^2,$$


because of $\gcd(|P(ix)|^2, |Q(ix)|^2) = (1 + 3x^2 + x^4)^2$. The problem can be additionally simplified by taking (see Lemma 1): $p(t) = 9 + 16t + 8t^2 + t^3$, $q(t) = 7 + 16t + 8t^2 + t^3$, $h(t) = 2t(p(t)q'(t) - q(t)p'(t)) = 2t(16 + 16t + 3t^2)$, and

$$R(t) = 2 \frac{h(t)}{p(t)q(t)} = \frac{4t(16 + 16t + 3t^2)}{(7 + 16t + 8t^2 + t^3)(9 + 16t + 8t^2 + t^3)}.$$

Fig. 1. The function $R(x^2)$

Thus, $R(t) \in \mathcal{R}[3, 6]$. The behavior of the function $R(x^2)$ is presented in Fig. 1. Its values for $x = \pm 5, \pm 10, \pm 15, \pm 20$ are 0.000518, 0.0000108, 1.005×10^{-6} , 1.826×10^{-7} , respectively.

However, after DE transformation $x = u(t) = \sinh((\pi/2) \sinh t)$, the integrand $R(u(t)^2)u'(t)$ decays double exponentially (see Fig. 2). For example, its values for $t = \pm 1, \pm 2, \pm 3 \pm 4$ are 0.04298, 9.654×10^{-10} , 4.102×10^{-31} , 1.357×10^{-89} , respectively.

Fig. 2. The function $R(u(t)^2)u'(t)$

Taking the bounds in the integral as $a = -3$ and $b = 3$ (corresp. value of integrand 4.102×10^{-31}), for $N = 10(10)70$ we get the trapezoidal approximations $I_h^{(N+1)}$. Table 1 shows these approximations, together with the relative errors. In each entry of the second column the first digit in error is underlined>. In the third column numbers in parentheses indicate decimal exponents, for example $1.40(-2) = 1.40 \times 10^{-2}$.

Table 1: Numerical approximations $I_h^{(N+1)}$ and the corresponding relative errors for $N = 10(10)100$

N	$I_h^{(N+1)}$	e_{rel}
10	0.385796	1.40(-2)
20	0.38049435	4.33(-5)
30	0.38047789486	7.92(-8)
40	0.38047786477677	1.25(-10)
50	0.3804778647293509	2.21(-13)
60	0.3804778647292668368	3.98(-16)
70	0.380477864729266856998	6.90(-19)
80	0.380477864729266854378032	1.20(-21)
90	0.38047786472926685437346027	2.11(-24)
100	0.380477864729266854373452238	3.70(-27)

The exact value (to 33 significant digits), as determined by the method in the next section, is 0.38047786472926685437345222424304. The corresponding exact value of (6) is $ef(G, Z) = 0.121109865817424581769007\dots$

4. Transformation to the finite interval and Gaussian formulae

In this section we propose another transformation $x = -t/\sqrt{1-t^2}$ (cf. [88]) in order to reduce (9) to the following integral over the finite interval $(-1, 1)$,

$$\int_{-\infty}^{+\infty} R(x^2) dx = \int_{-1}^1 R\left(\frac{t^2}{1-t^2}\right) \frac{dt}{(1-t^2)^{3/2}}, \quad (13)$$

where $R(t)$ is defined in (8). This suggests us to apply some of Gaussian formulas for numerical calculation of (13). Namely, the Gaussian quadrature rule with respect to the Gegenbauer weight $w^\lambda(t) = (1-t^2)^{\lambda-1/2}$, $\lambda > -1/2$,

$$\int_{-1}^1 \varphi(t) w^\lambda(t) dt = \sum_{k=1}^N A_k^\lambda \varphi(\tau_k^\lambda) + R_N(\varphi), \quad (14)$$

could be appropriate for this purpose. The nodes τ_k^λ , $k = 1, \dots, N$, are zeros of the Gegenbauer polynomial $C_N^\lambda(t)$ of degree N , and the weights A_k^λ , $k = 1, \dots, N$, are the corresponding Christoffel numbers (cf. [84, Chap. 5]). They can be calculated in an efficient way by using the MATHEMATICA Package “OrthogonalPolynomials” [11].

Example 2 Consider again the integral from Example 1. In this case, the integral (13), written as a weighted integral with respect to the Gegenbauer weight $w^\lambda(t)$, becomes

$$\int_{-1}^1 \frac{4t^2 (3t^4 - 16t^2 + 16) (1 - t^2)^{2-\lambda}}{(3t^4 - 11t^2 + 9) (2t^6 - 3t^4 - 5t^2 + 7)} w^\lambda(t) dt.$$

The complete integrand $R(t^2/(1-t^2))(1-t^2)^{-3/2}$ is presented in Fig. 3. Applying the corresponding Gaussian formula (14) for $\lambda = 0, 1/2, 1, 3/2, 2$ to the previous integral we obtain results with the relative errors presented in Table 2. Note that this parameter λ must be such that $-1/2 < \lambda \leq 2$.




Fig. 3. The function $R(t^2/(1-t^2))(1-t^2)^{-3/2}$ in Example 2

Table 2. Relative errors in Gauss-Gegenbauer quadrature sums for some selected values of λ

N	$\lambda = 0$	$\lambda = 1/2$	$\lambda = 1$	$\lambda = 3/2$	$\lambda = 2$
10	9.99(-4)	1.18(-3)	4.28(-4)	1.24(-3)	2.75(-4)
20	1.64(-7)	2.77(-5)	6.72(-8)	3.75(-5)	3.42(-8)
30	2.06(-11)	3.39(-6)	8.34(-12)	5.16(-6)	3.90(-12)
40	2.33(-15)	7.95(-7)	9.35(-16)	1.27(-6)	4.20(-16)
50	2.48(-19)	2.60(-7)	9.91(-20)	4.27(-7)	4.33(-20)
60	2.54(-23)	1.04(-7)	1.01(-23)	1.75(-7)	4.35(-24)
70	2.54(-27)	4.84(-8)	1.01(-27)	8.19(-8)	4.28(-28)
80	2.49(-31)	2.48(-8)	9.89(-32)	4.25(-8)	4.16(-32)
90	2.41(-35)	1.38(-8)	9.56(-36)	2.38(-8)	3.99(-36)
100	2.31(-39)	8.16(-9)	9.14(-40)	1.41(-8)	3.79(-40)

As we can see, the convergence is slow only if the integrand has an irrational factor. In these cases it is the factor “ $\sqrt{1-t^2}$ ” which appears

for $\lambda = 1/2$ (Gauss-Legendre quadrature) and $\lambda = 3/2$. In other cases this factor is included in the weight function and the corresponding function $\varphi(t)$ is a pure rational function. Regarding this fact we prefer to use Gauss-Chebyshev quadrature formulas (for $\lambda = 0$ and $\lambda = 1$), because of their simplicity.

In a general case, the rational function $R(t)$ belongs to $\mathcal{R}[m, 2n]$, where $0 \leq m \leq 2n - 1$ (see Lemma 1), and therefore it has the following form

$$R(t) = \frac{\sum_{k=0}^m r_k t^k}{\sum_{k=0}^{2n} s_k t^k}.$$

Regarding (13), we have

$$\int_{-\infty}^{+\infty} R(x^2) dx = \int_{-1}^1 \frac{\sum_{k=0}^m r_k t^{2k} (1-t^2)^{m-k}}{\sum_{k=0}^{2n} s_k t^{2k} (1-t^2)^{2n-k}} (1-t^2)^\ell \cdot \frac{dt}{\sqrt{1-t^2}},$$

where $\ell = 2n - m - 1 \geq 0$. Thus, the Gauss-Chebyshev quadrature of the first kind ($\lambda = 0$) can be always applied to the integral (13). However, the corresponding Gauss-Chebyshev quadrature of the second kind ($\lambda = 1$) can be applied if $m \leq 2n - 2$.

Now, we derive explicit expressions for these Gaussian quadrature sums

$$\int_{-\infty}^{+\infty} R(x^2) dx \approx S_N^\lambda(R) \quad (\lambda = 0, 1). \quad (15)$$

Number of functional evaluations in these sums is reduced to $N/2$.

Theorem 1. *Let $N \in \mathbb{N}$, $R(t) \in \mathcal{R}[m, 2n]$, $0 \leq m \leq 2n - 1$, and $\xi_k = \cot^2 \frac{(2k-1)\pi}{2N}$, $k = 1, \dots, [N/2]$. Then*

$$S_N^T(R) = \frac{2\pi}{N} \sum_{k=1}^{[N/2]} (1 + \xi_k) R(\xi_k) + \varepsilon_N \frac{\pi}{N} R(0), \quad (16)$$

where $\varepsilon_N = 0$ if N is even, and $\varepsilon_N = 1$ if N is odd.

P r o o f: Let $\lambda = 0$ and $\varphi(t)$ be defined as

$$\varphi(t) = \frac{1}{1-t^2} R\left(\frac{t^2}{1-t^2}\right).$$

Then for the N -point Gauss-Chebyshev quadrature sum of the first kind in (14), with the nodes $\tau_k = \tau_k^0 = \cos \theta_k$, where $\theta_k = \frac{(2k-1)\pi}{2N}$, $k = 1, \dots, N$, all weight coefficients are equal, i.e., $A_k = A_k^0 = \pi/N$ (cf. [86, p. 174]). Therefore,

$$S_N^0(R) = \frac{\pi}{N} \sum_{k=1}^N \frac{1}{1-\cos^2 \theta_k} R\left(\frac{\cos^2 \theta_k}{1-\cos^2 \theta_k}\right) = \frac{\pi}{N} \sum_{k=1}^N \frac{1}{\sin^2 \theta_k} R(\cot^2 \theta_k),$$

i.e.,

$$S_N^0(R) = \frac{\pi}{N} \sum_{k=1}^N (1 + \cot^2 \theta_k) R(\cot^2 \theta_k),$$

which reduces to (16). \square

Theorem 2. Let $N \in \mathbb{N}$, $R(t) \in \mathcal{R}[m, 2n]$, $0 \leq m \leq 2n - 2$, and $\eta_k = \cot^2 \frac{k\pi}{N+1}$, $k = 1, \dots, [N/2]$. Then

$$S_N^1(R) = \frac{2\pi}{N+1} \sum_{k=1}^{[N/2]} (1 + \eta_k) R(\eta_k) + \varepsilon_N \frac{\pi}{N+1} R(0), \quad (17)$$

where $\varepsilon_N = 0$ if N is even, and $\varepsilon_N = 1$ if N is odd.

P r o o f: In this case $\lambda = 1$ and

$$\varphi(t) = \frac{1}{(1-t^2)^2} R\left(\frac{t^2}{1-t^2}\right).$$

Nodes of the corresponding N -point quadrature are zeros of the Chebyshev polynomial of the second kind $U_N(t) = \sin[(N+1)\arccos t]/\sqrt{1-t^2}$, i.e., $\tau_k = \tau_k^1 = \cos \theta_k$, where $\theta_k = \frac{k\pi}{N+1}$, $k = 1, \dots, N$, and the weight coefficients are $A_k = A_k^1 = \frac{\pi}{N+1} \sin^2 \theta_k$, $k = 1, \dots, N$ (cf. [86, p. 174]). Therefore,

$$S_N^1(R) = \frac{\pi}{N+1} \sum_{k=1}^N \frac{\sin^2 \theta_k}{(1-\cos^2 \theta_k)^2} R\left(\frac{\cos^2 \theta_k}{1-\cos^2 \theta_k}\right)$$

reduces to (17). □

The obtained formulas (16) and (17) are very simple for implementing and using in integration. For example, the function $R(t)$ from Examples 1 and 2 belongs to $\mathcal{R}[3, 6]$ and Theorem 2 can be applied. The quadrature sums $S_1^U(R)$ for $N = 10(10)100$ are presented in Table 3.

Table 3. Quadrature sums $S_N^1(R)$ for $N = 10(10)100$ values of λ

N	Quadrature sum S_N^1
10	0.3806407
20	0.38047789027
30	0.38047786473243
40	0.380477864729267041
50	0.38047786472926668547505
60	0.380477864729266685437349077
70	0.3804778647292666854373452228086
80	0.38047786472926668543734522242434194
90	0.380477864729266685437345222424304306513
100	0.3804778647292666854373452224243043028756715

REFERENCES

- [1] J. Aihara, *Resonance energies of benzenoid hydrocarbons*, J. Am. Chem. Soc. **99** (1977) 2048–2053.
- [2] J. Aihara, *Circuit resonance energy: A key quantity that links energetic and magnetic criteria of aromaticity*, J. Am. Chem. Soc. **128** (2006) 2873–2879.
- [3] A. T. Balaban, J. Đurđević, I. Gutman, *Comments on π -electron conjugation in the five-membered ring of benzo-derivatives of corannulene*, Polyc. Arom. Comp. **29** (2009) 185–205.
- [4] A. T. Balaban, J. Đurđević, I. Gutman, S. Jeremić, S. Radenković, *Correlations between local aromaticity indices of bipartite conjugated hydrocarbons*, J. Phys. Chem. **A 114** (2010) 5870–5877.
- [5] A. T. Balaban, I. Gutman, S. Jeremić, J. Đurđević, *Effect of benzo-annulation on cyclic conjugation*, Monatsh. Chem. **142** (2011) 53–57.
- [6] G. Boros, V. H. Moll, *Landen transformations and the integration of rational functions*, Math. Comp. **71** (2002), 649–668.
- [7] S. Bosanac, I. Gutman, *Effect of a ring on the stability of polycyclic conjugated molecules*, Z. Naturforsch. **32a** (1977) 10–12.
- [8] M. Bronstein, *Symbolic Integration I. Transcendental functions*, Second Edition, Algorithms and Computation in Mathematics, Vol. 1, Springer-Verlag, Berlin – Heidelberg, 2005.
- [9] C. A. Coulson, *On the calculation of the energy in unsaturated hydrocarbon molecules*, Proc. Cambridge Phil. Soc. **36** (1940) 201–203.

- [10] C. A. Coulson, J. Jacobs, *Conjugation across a single bond*, J. Chem. Soc. (1949) 2805–2812.
- [11] A. S. Cvetković, G. V. Milovanović, *The Mathematica Package “OrthogonalPolynomials”*, Facta Univ. Ser. Math. Inform. **19** (2004), 17–36.
- [12] J. Đurđević, B. Furtula, I. Gutman, R. Kovačević, S. Stanković, N. Turković, *Cyclic conjugation in annelated perylenes*, in: I. Gutman (Ed.), *Mathematical Methods in Chemistry*, Prijepolje Museum, Prijepolje, 2006, pp. 101–117.
- [13] J. Đurđević, B. Furtula, I. Gutman, S. Radenković, S. Stanković, *Comparative study of cyclic conjugation in tribenzoperylene isomers*, in: A. Graovac, I. Gutman, D. Vukičević (Eds.), *Mathematical Methods and Modelling for Students of Chemistry and Biology*, Hum, Zagreb, 2009, pp. 29–39.
- [14] J. Đurđević, I. Gutman, J. Terzić, A. T. Balaban, *Cyclic conjugation in fluoranthene and its benzo-derivatives. Part 1. Catacondensed systems*, Polyc. Arom. Comp. **29** (2009) 90–102.
- [15] J. Đurđević, S. Radenković, I. Gutman, S. Marković, *Testing the PCP-rule*, Monatsh. Chem. **140** (2009) 1305–1309.
- [16] B. Furtula, I. Gutman, S. Jeremić, S. Radenković, *Effect of a ring on cyclic conjugation in another ring: Applications to acenaphthylene-type polycyclic conjugated molecules*, J. Serb. Chem. Soc. **75** (2010) 83–90.
- [17] I. Gutman, *Proof of the Hückel rule*, Chem. Phys. Lett. **46** (1977) 169–171.
- [18] I. Gutman, *Electronic properties of Möbius systems* Z. Naturforsch. **33a** (1978) 214–216.
- [19] I. Gutman, *Topological studies on heteroconjugated molecules. Alternant systems with one heteroatom*, Theor. Chim. Acta **50** (1979) 287–297.
- [20] I. Gutman, *Effect of cycles on total π -electron energy of alternant conjugated hydrocarbons*, J. Chem. Soc. Faraday Trans. II **75** (1979) 799–805.
- [21] I. Gutman, *Two theorems on topological resonance energy*, J. Serb. Chem. Soc. **44** (1979) 173–178.
- [22] I. Gutman, *Effect of cycles on topological resonance energy*, Croat. Chem. Acta **53** (1980) 581–586.
- [23] I. Gutman, *On cyclic conjugation*, Theor. Chim. Acta **66** (1984) 43–49.
- [24] I. Gutman, *Overlooked relations in the theory of cyclic conjugation*, Chem. Phys. Lett. **117** (1985) 614–616.
- [25] I. Gutman, *Cyclic conjugation in porphin*, Croat. Chem. Acta **58** (1985) 359–369.
- [26] I. Gutman, *Cyclic conjugation in fully benzenoid hydrocarbons*, Rep. Mol. Theory **1** (1990) 115–119.
- [27] I. Gutman, *Dependence of physico-chemical properties of substances on molecular structure: The example of total π -electron energy*, Glas Acad. Serbe Sci. Arts **362** (1990) 83–91 (in Serbian).
- [28] I. Gutman, *Topological studies on heteroconjugated molecules. VI. Alternant systems with two heteroatoms*, Z. Naturforsch. **45a** (1990) 1085–1089.

- [29] I. Gutman, *Violations of the Hückel ($4n + 2$) rule*, J. Chin. Chem. Soc. **40** (1993) 7–10.
- [30] I. Gutman, *Hückel rule in catacondensed benzenoid hydrocarbons*, MATCH Commun. Math. Comput. Chem. **29** (1993) 51–60.
- [31] I. Gutman, *Cyclic conjugation in benzenoid hydrocarbons*, Bull. Acad. Serbe Sci. Arts **106** (1993) 37–50.
- [32] I. Gutman, *Cyclic conjugation in phenylenes*, S. Afr. J. Chem. **47** (1994) 53–55.
- [33] I. Gutman, *Cyclic conjugation in antikekulene and its homologues*, Rev. Roum. Chim. **39** (1994) 943–947.
- [34] I. Gutman, *Cyclic conjugation in dianions: effect of cycles on the thermodynamic stability of polycyclic conjugated dianions*, J. Mol. Struct. (Theochem) **428** (1998) 241–246.
- [35] I. Gutman, *On the oxidation of dibenzo[fg,ij]pentaphene and, dinaphtho[2,1,8,7-defg:2',1',8',7'-ijkl]pentaphene dianions*, J. Serb. Chem. Soc. **64** (1999) 563–570.
- [36] I. Gutman, *Cyclic conjugation energy effects in polycyclic π -electron systems*, Monatsh. Chem. **136** (2005) 1055–1069.
- [37] I. Gutman, *What chemists could not see without mathematics – Dependence of total π -electron energy on molecular structure*, Kragujevac J. Sci. **27** (2005) 57–66.
- [38] I. Gutman, *What chemists could not see without mathematics – Dependence of total π -electron energy on molecular structure*, Xjenza (Malta) **10** (2005) 3–7.
- [39] I. Gutman, *Theory of the PCP effect and related phenomena*, J. Math. Chem. **47** (2010) 1309–1312.
- [40] I. Gutman, *On a class of integrals encountered in theoretical chemistry*, Int. J. Chem. Model. **2** (2010) 335–341.
- [41] I. Gutman, P. U. Biedermann, V. Ivanov–Petrović, I. Agranat, *Cyclic conjugation effects in cyclacenes*, Polyc. Arom. Comp. **8** (1996) 189–202.
- [42] I. Gutman, S. Bosanac, *Topological studies on heteroconjugated molecules. The stability of alternant systems with one heteroatom*, Chem. Phys. Lett. **43** (1976) 371–373.
- [43] I. Gutman, S. Bosanac, *Quantitative approach to Hückel rule. The relations between the cycles of a molecular graph and the thermodynamic stability of a conjugated molecule*, Tetrahedron **33** (1977) 1809–1812.
- [44] I. Gutman, S. Bosanac, *Quantitative testing of the perimeter rule*, J. Serb. Chem. Soc. **42** (1977) 499–502.
- [45] I. Gutman, S. J. Cyvin, V. Petrović, A. Teodorović, *Fully-naphthalenoid hydrocarbons and their conjugation modes*, Polyc. Arom. Comp. **4** (1994) 183–189.
- [46] I. Gutman, J. Đurđević, *On π -electron conjugation in the five-membered ring of fluoranthene-type benzenoid hydrocarbons*, J. Serb. Chem. Soc. **74** (2009) 765–771.
- [47] I. Gutman, J. Đurđević, A. T. Balaban, *A regularity for cyclic conjugation in acenaphthylene, fluoranthene and their congeners*, Polyc. Arom. Comp. **29** (2009) 3–11.
- [48] I. Gutman, J. Đurđević, B. Furtula, B. Milivojević, *Cyclic conjugation in mono- and dicyclopenta-derivatives of anthracene and phenanthrene*, Indian J. Chem. **47A** (2008) 803–807.

- [49] I. Gutman, B. Furtula, *Cyclic conjugation in pyracylene*, Polyc. Arom. Comp. **28** (2008) 136–142.
- [50] I. Gutman, B. Furtula, J. Đurđević, R. Kovačević, S. Stanković, *Annulated perylenes: Benzenoid molecules violating the Kekulé-structure-based cyclic conjugation models*, J. Serb. Chem. Soc. **70** (2005) 1023–1031.
- [51] I. Gutman, V. Ivanov–Petrović, *Unusual modes of cyclic conjugation in phenylenes*, Bull. Chem. Technol. Maced. **16** (1997) 91–96.
- [52] I. Gutman, V. Ivanov–Petrović, *Clar theory and phenylenes*, J. Mol. Struct. (Theochem) **389** (1997) 227–232.
- [53] I. Gutman, V. Ivanov–Petrović, J. R. Dias, *Cyclic conjugation in total resonant sextet benzenoid hydrocarbons*, Polyc. Arom. Comp. **18** (2000) 221–229.
- [54] I. Gutman, V. Ivanov–Petrović, S. Pogodin, *Unusual conjugation pattern in a class of benzenoid hydrocarbons*, Indian J. Chem. **35A** (1996) 87–92.
- [55] I. Gutman, S. Jeremić, V. Petrović, *Extending the PCP rule*, Indian J. Chem. **48A** (2009) 658–662.
- [56] I. Gutman, S. L. Lee, *Cyclic conjugation in circulenes*, Bull. Inst. Chem. Acad. Sin. **40** (1993) 47–54.
- [57] I. Gutman, Z. Marković, I. Juranić, *A Clar-type regularity for dianions of benzenoid hydrocarbons*, Polyc. Arom. Comp. **13** (1999) 33–40.
- [58] I. Gutman, M. Mateljević, *Note on the Coulson integral formula*, J. Math. Chem. **39** (2006) 259–266.
- [59] I. Gutman, M. Milun, N. Trinajstić, *Topological definition of delocalisation energy*, MATCH Commun. Math. Comput. Chem. **1** (1975) 171–175.
- [60] I. Gutman, M. Milun, N. Trinajstić, *Graph theory and molecular orbitals. 19. Non-parametric resonance energies of arbitrary conjugated systems*, J. Am. Chem. Soc. **99** (1977) 1692–1704.
- [61] I. Gutman, V. Petrović, *Cyclic conjugation in benzo-annulated polyacenes*, Indian J. Chem. **31A** (1992) 647–650.
- [62] I. Gutman, V. Petrović, *Cyclic conjugation in linear polyacenes*, J. Serb. Chem. Soc. **57** (1992) 495–501.
- [63] I. Gutman, V. Petrović, *Cyclic conjugation in zig-zag fibonacenes*, Rev. Roum. Chim. **38** (1993) 439–445.
- [64] I. Gutman, V. Petrović, *Theoretical study of cyclic conjugation in phenes: Some hitherto unnoticed features*, Monatsh. Chem. **126** (1995) 1179–1185.
- [65] I. Gutman, V. Petrović, B. Mohar, *Cyclic conjugation effects: individual, collective and overall*, Chem. Phys. Lett. **203** (1993) 378–382.
- [66] I. Gutman, O. E. Polansky, *Cyclic conjugation and the Hückel molecular orbital model*, Theor. Chim. Acta **60** (1981) 203–226.
- [67] I. Gutman, O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, Berlin, 1986.
- [68] I. Gutman, S. Radenković, W. Linert, *Pairwise energy effect of cyclic conjugation in benzo-annulated perylenes*, Monatsh. Chem. **141** (2010) 401–407.

- [69] I. Gutman, M. Stanković, *Violation of the Hückel rule in alternant nonbenzenoid hydrocarbons*, J. Serb. Chem. Soc. **58** (1993) 897–903.
- [70] I. Gutman, M. Stanković, *Cyclic conjugation in non-Kekulean benzenoid molecules*, Coll. Sci. Pap. Fac. Sci. Kragujevac **15** (1994) 97–104.
- [71] I. Gutman, M. Stanković, *On the Hückel $(4m + 2)$ rule in polycyclic alternant hydrocarbons*, J. Mol. Struct. (Theochem) **309** (1994) 301–304.
- [72] I. Gutman, S. Stanković, *Testing the Y-rule in Clar theory*, Polyc. Arom. Comp. **27** (2007) 425–436.
- [73] I. Gutman, S. Stanković, *Why is phenanthrene more stable than anthracene?*, Maced. J. Chem. Chem. Engin. **26** (2007) 111–114.
- [74] I. Gutman, S. Stanković, J. Đurđević, B. Furtula, *On the cycle-dependence of topological resonance energy*, J. Chem. Inf. Model. **47** (2007) 776–781.
- [75] I. Gutman, S. Stanković, *Complete analysis of cyclic conjugation energy effects: Comparison of, benzo[a]pyrene and benzo[b]pyrene*, Bull. Acad. Serbe Sci. Arts **135** (2008) 29–36.
- [76] I. Gutman, S. Stanković, R. Kovačević, J. Đurđević, B. Furtula, *Anomalous cyclic conjugation in benzenoid molecules with a small number of Kekulé structures*, Indian J. Chem. **44A** (2005) 1751–1755.
- [77] I. Gutman, Ž. Tomović, *On cyclic conjugation of the members of the pyrene/peropyrene series and their formally π -localized derivatives*, Bull. Chem. Technol. Maced. **20** (2001) 33–37.
- [78] I. Gutman, Ž. Tomović, *Cyclic conjugation in terminally bent and branched phenylenes*, Indian J. Chem. **40A** (2001) 678–681.
- [79] I. Gutman, N. Turković, J. Jovičić, *Cyclic conjugation in benzo-annelated perylenes: How empty is the “empty” ring ?*, Monatsh. Chem. **135** (2004) 1389–1394.
- [80] M. Iri, S. Moriguti, Y. Takasawa, *On certain quadrature formula*, J. Comput. Appl. Math. **17** (1987), 3–20.
- [81] S. Jeremić, S. Radenković, I. Gutman, *Cyclic conjugation in benzo-annelated coronenes*, Maced. J. Chem. Chem. Engin. **29** (2010) 63–69.
- [82] S. Jeremić, S. Radenković, I. Gutman, *Cyclic conjugation in benzo-annelated triphenylenes*, J. Serb. Chem. Soc. **75** (2010) 943–950.
- [83] D. Manna, V. H. Moll, *Rational Landen transformations on \mathbb{R}* , Math. Comp. **76** (2007), 2023–2043.
- [84] G. Mastraianni, G. V. Milovanović, *Interpolation Processes – Basic Theory and Applications*, Springer Monographs in Mathematics, Springer Verlag, Berlin – Heidelberg – New York, 2008.
- [85] M. Mateljević, I. Gutman, *Note on the Coulson and Coulson–Jacobs integral formulas*, MATCH Commun. Math. Comput. Chem. **59** (2008) 257–268.
- [86] G. V. Milovanović, *Numerical Analysis, Part II*, Naučna knjiga, Beograd, 1988 (Serbian).
- [87] G. V. Milovanović, *Expansions of the Kurepa function*, Publ. Inst. Math. (Beograd) (N.S.) **57 (71)** (1995), 81–90.

- [88] G. Monegato, L. Scuderi, *Quadrature rules for unbounded intervals and their application to integral equations*, In: Approximation and Computation: In Honor of Gradimir V. Milovanović (W. Gautschi, G. Mastroianni, Th.M. Rassias, eds.), Springer Optimization and its Applications, Vol. 42, Springer, New York, 2011, pp. 185–208.
- [89] M. Mori, *Quadrature formulas obtained by variable transformation and DE-rule*, J. Comput. Appl. Math. **12&13** (1985), 119–130.
- [90] R. Ponec, S. Fias, S. Van Damme, P. Bultinck, I. Gutman, S. Stanković, *The close relation between cyclic delocalization, energy effects of cycles and aromaticity*, Coll. Czech. Chem. Commun. **74** (2009) 147–166.
- [91] S. Radenković, J. Đurđević, I. Gutman, *Quantitative study of the PCP effect*, Chem. Phys. Lett. **475** (2009) 289–292.
- [92] S. Radenković, W. Linert, I. Gutman, S. Jeremić, *Pairwise energy effects of rings in benzo-annelated perylenes*, Indian J. Chem. **48A** (2009) 1657–1661.
- [93] H. Takahasi, M. Mori, *Error estimation in the numerical integration of analytic functions*, Report Computer Centre University of Tokyo **3** (1970), 41–108.
- [94] H. Takahasi, M. Mori, *Quadrature formulas obtained by variable transformation*, Numer. Math. **21** (1973), 206–219.
- [95] J. Waldvogel, *Towards a general error theory of the trapezoidal rule*, In: Approximation and Computation: In Honor of Gradimir V. Milovanović (W. Gautschi, G. Mastroianni, Th.M. Rassias, eds.), Springer Optimization and its Applications, Vol. 42, Springer, New York, 2011, pp. 267–282.

Faculty of Science
University of Kragujevac
P. O. Box 60
34000 Kragujevac
Serbia
gutman@kg.ac.rs

Megatrend University
Faculty of Computer Sciences
Bulevar umetnosti 29
11070 Novi Beograd
Serbia
gvm@megatrend.edu.rs