

Graph Eigenvalues in Combinatorial Optimization

Domingos Moreira Cardoso¹

¹CIDMA - DMAT - Universidade de Aveiro, Portugal

16th ILAS Conference - Minisymposia on Spectral Graph Theory,
June 21-25, 2010, Pisa, Italy

Outline

Classical results

Spectral bounds on the chromatic and stability number
Maximum cut and quadratic assignment problems

Recent results

Maximum size induced k -regular subgraphs
Perfect Matchings and Hamiltonian graphs

(k, τ) -regular sets

Definition and particular cases
Combinatorial and algebraic properties
Main and non-main eigenvalue techniques

Bibliography

Graph eigenvalues and the chromatic number

- ▶ In some sense, we may say that the graph eigenvalues encode the combinatorial structure of the graph.

Graph eigenvalues and the chromatic number

- ▶ In some sense, we may say that the graph eigenvalues encode the combinatorial structure of the graph. This allows several applications in Combinatorial Optimization.

Graph eigenvalues and the chromatic number

- ▶ In some sense, we may say that the graph eigenvalues encode the combinatorial structure of the graph. This allows several applications in Combinatorial Optimization.
- ▶ Considering a graph G , the inequalities

$$1 + \frac{\lambda_1(G)}{-\lambda_n(G)} \leq \chi(G) \leq \lambda_1(G) + 1.$$

are the earliest graph spectra applications in combinatorial optimization.

Graph eigenvalues and the chromatic number

- ▶ In some sense, we may say that the graph eigenvalues encode the combinatorial structure of the graph. This allows several applications in Combinatorial Optimization.
- ▶ Considering a graph G , the inequalities

$$1 + \frac{\lambda_1(G)}{-\lambda_n(G)} \leq \chi(G) \leq \lambda_1(G) + 1.$$

are the earliest graph spectra applications in combinatorial optimization.

- ▶ The left inequality was obtained by Hoffman in **1970** and the right by Wilf in **1967**.

A lower bound on the stability number

On that time, based on the quadratic program of [Motzkin and Straus](#) [[Motzkin and Straus, 1965](#)], [Wilf](#) also proposed a spectral lower bound on the stability number of a graph.

A lower bound on the stability number

On that time, based on the quadratic program of [Motzkin and Straus \[Motzkin and Straus, 1965\]](#), [Wilf](#) also proposed a spectral lower bound on the stability number of a graph.

Theorem [Wilf, 1967]

Let G be a graph with at least one edge. Then

$$\alpha(G) \geq \frac{s^2}{s^2 - \lambda_1(\overline{G})},$$

where s is the sum of the entries of the normalized eigenvector corresponding to $\lambda_1(\overline{G})$ (the index of the complement of G).

Another lower bound on the chromatic number

Taking into account that

$$\chi(G) \geq \omega(G) = \alpha(\overline{G}) \geq \frac{s^2}{s^2 - \lambda_1(G)},$$

$s^2 \leq n$ and $\frac{s^2}{s^2 - \lambda_1(G)}$ is a decreasing function on s^2 , the **Cvetković's bound** on chromatic number can be deduced.

Another lower bound on the chromatic number

Taking into account that

$$\chi(G) \geq \omega(G) = \alpha(\overline{G}) \geq \frac{s^2}{s^2 - \lambda_1(G)},$$

$s^2 \leq n$ and $\frac{s^2}{s^2 - \lambda_1(G)}$ is a decreasing function on s^2 , the **Cvetković's bound** on chromatic number can be deduced.

Theorem [Cvetković, 1972]

If $G \neq K_n$ is a graph with n vertices, then

$$\chi(G) \geq \frac{n}{n - \lambda_1(G)}.$$

Upper bounds on the stability number

Theorem [Ryacek and Sciermeyer, 1995]

If G is a claw-free graph with n vertices and at least one edge, then $\alpha(G) \leq \frac{2n}{2+\delta(G)}$.

Upper bounds on the stability number

Theorem [Ryacek and Sciermeyer, 1995]

If G is a claw-free graph with n vertices and at least one edge, then $\alpha(G) \leq \frac{2n}{2+\delta(G)}$.

As a consequence (since $\alpha(G)\chi(G) \geq n$), if G is k -regular and claw-free, then $\chi(G) \geq \frac{k+2}{2}$.

Upper bounds on the stability number

Theorem [Ryacek and Sciermeyer, 1995]

If G is a claw-free graph with n vertices and at least one edge, then $\alpha(G) \leq \frac{2n}{2+\delta(G)}$.

As a consequence (since $\alpha(G)\chi(G) \geq n$), if G is k -regular and claw-free, then $\chi(G) \geq \frac{k+2}{2}$.

Theorem [Cvetković, 1973]

Let $|\sigma^-(G)|$, $|\sigma^0(G)|$, $|\sigma^+(G)|$ the number of eigenvalues of the graph G smaller than, equal to, and greater than zero, respectively. Then

$$\alpha(G) \leq |\sigma^0(G)| + \min\{|\sigma^-(G)|, |\sigma^+(G)|\}.$$

The maximum cut problem

The maximum cut (or max-cut) problem is defined as follows.

The maximum cut problem

The maximum cut (or max-cut) problem is defined as follows.

Determine a vertex subset $S \subset V(G)$ for which the weight

$w(\partial(S)) = \sum_{e \in \partial(S)} w(e)$ is maximum, that is,

$$\max_{S \subset V(G)} w(\partial(S)). \quad (1)$$

The maximum cut problem

The maximum cut (or max-cut) problem is defined as follows. Determine a vertex subset $S \subset V(G)$ for which the weight $w(\partial(S)) = \sum_{e \in \partial(S)} w(e)$ is maximum, that is,

$$\max_{S \subset V(G)} w(\partial(S)). \quad (1)$$

This problem can be formulated as the following integer quadratic program:

$$\begin{aligned} mc(G, w) \quad = \quad & \text{Maximize} \quad \frac{1}{2} \sum_{ij \in E(G)} w_{ij} (1 - x_i x_j) \\ & \text{subject to } x_i \in \{1, -1\} \quad \forall i \in V(G). \end{aligned}$$

Upper bound on the maximum cut problem

Theorem [Mohar and Poljak, 1990]

Let $L_{(G,w)}$ be the Laplacian matrix of the weighted graph (G, w) .

Then

$$mc(G, w) \leq \frac{n}{4} \lambda_1(L_{(G,w)}).$$

Upper bound on the maximum cut problem

Theorem [Mohar and Poljak, 1990]

Let $L_{(G,w)}$ be the Laplacian matrix of the weighted graph (G, w) .
Then

$$mc(G, w) \leq \frac{n}{4} \lambda_1(L_{(G,w)}).$$

Theorem [Delorme and Poljak, 1990]

Let $L_{(G,w)}$ be the Laplacian matrix of the weighted graph (G, w) . Then

$$mc(G, w) \leq \min_u \frac{n}{4} \lambda_1(L_{(G,w)} + \text{diag}(u)),$$

where the minimum is obtained among overall $u \in \mathbb{R}^n$ such that $\sum_i u_i = 0$.

- └ Classical results
 - └ Maximum cut and quadratic assignment problems

The quadratic assignment problem

- ▶ The *quadratic assignment* problem is one the most general combinatorial optimization models.

The quadratic assignment problem

- ▶ The *quadratic assignment* problem is one the most general combinatorial optimization models. Many other hard combinatorial optimization problems (as it is te case of the location decision problem) are particular cases of the quadratic assignment problem.

The quadratic assignment problem

- ▶ The *quadratic assignment* problem is one the most general combinatorial optimization models. Many other hard combinatorial optimization problems (as it is the case of the location decision problem) are particular cases of the quadratic assignment problem.
- ▶ The **quadratic assignment problem** (QAP) can be defined as follows: given the set $\mathbf{N} = \{1, \dots, n\}$ and three $n \times n$ matrices $\mathbf{A} = (a_{ik})$, $\mathbf{B} = (b_{jl})$ and $\mathbf{C} = (c_{ij})$, find a permutation π of the set \mathbf{N} which minimizes

$$\sum_{i=1}^n c_{i\pi(i)} + \sum_{i=1}^n \sum_{j=1}^n a_{ij} b_{\pi(i)\pi(j)}.$$

The quadratic assignment problem (alternative formulation)

- Equivalently, QAP consists on finding a $n \times n$ permutation matrix P which minimizes the trace

$$\min_{P \in \Pi} \text{tr}(C + APB^t)P^t,$$

where Π denotes the set of permutation matrices.

The quadratic assignment problem (alternative formulation)

- Equivalently, QAP consists on finding a $n \times n$ permutation matrix P which minimizes the trace

$$\min_{P \in \Pi} \text{tr}(C + APB^t)P^t,$$

where Π denotes the set of permutation matrices.

► Theorem [Finke, Burkard and Rendl, 1987]

Let A and B be symmetric $n \times n$ matrices with eigenvalues $\lambda_1 \geq \dots \geq \lambda_n$ and $\mu_1 \geq \dots \geq \mu_n$, respectively. Then

$$\min_{\pi} \sum_{i=1}^n \sum_{j=1}^n a_{ij} b_{\pi(i)\pi(j)} \geq \sum_{i=1}^n \lambda_i \mu_{n-i+1},$$

where the minimum is taken over all permutations π of $\{1, \dots, n\}$.

Upper bound on the size of a vertex-set inducing subgraph with mean degree k

- Using a quadratic programming technique jointly with the main angles of G , as proved in (C and Rowlinson, 2010), the size of a vertex subset $S \subset V(G)$ inducing a subgraph with mean degree k has the following upper bound:

$$|S| \leq \inf\{h_k^G(t) : t > -\lambda_n(G)\},$$

$$\text{where } h_k^G(t) = (k + t) \left(1 - \frac{P_G(t-1)}{(-1)^n P_G(-t)} \right).$$

Upper bound on the size of a vertex-set inducing subgraph with mean degree k

- ▶ Using a quadratic programming technique jointly with the main angles of G , as proved in (C and Rowlinson, 2010), the size of a vertex subset $S \subset V(G)$ inducing a subgraph with mean degree k has the following upper bound:

$$|S| \leq \inf\{h_k^G(t) : t > -\lambda_n(G)\},$$

where
$$h_k^G(t) = (k+t) \left(1 - \frac{P_G(t-1)}{(-1)^n P_G(-t)}\right).$$

- ▶ Considering $H_G(t) = \sum_{k=0}^{\infty} N_k t^k$, where N_k is the number of walks of length k in G (the walk-generating function of G), we may write $h_k^G(t) = (1 + \frac{k}{t}) H_G(-\frac{1}{t})$.

Induced matchings and dominating induced matchings

Lemma

Let G be a graph with an induced matching $M \subseteq E(G)$. Then

$$\triangleright |M| \leq \min\{|\sigma^+(G)|, |\sigma^-(G)|\};$$

Induced matchings and dominating induced matchings

Lemma

Let G be a graph with an induced matching $M \subseteq E(G)$. Then

$$\triangleright |M| \leq \min\{|\sigma^+(G)|, |\sigma^-(G)|\};$$

Theorem [C, Cerdeira, Delorme and Silva, 2008]

Let G be a graph. If $M \subseteq E(G)$ is a dominating induced matching, then $|M| \geq \frac{1}{2} \max\{|\sigma^+(G)|, |\sigma^-(G)|\}$.

Induced matchings and dominating induced matchings

Lemma

Let G be a graph with an induced matching $M \subseteq E(G)$. Then

$$\triangleright |M| \leq \min\{|\sigma^+(G)|, |\sigma^-(G)|\};$$

Theorem [C, Cerdeira, Delorme and Silva, 2008]

Let G be a graph. If $M \subseteq E(G)$ is a dominating induced matching, then $|M| \geq \frac{1}{2} \max\{|\sigma^+(G)|, |\sigma^-(G)|\}$.

Corollary

If M is a dominating induced matching of a graph G , then $\frac{1}{2} \max\{|\sigma^+(G)|, |\sigma^-(G)|\} \leq |M| \leq \min\{|\sigma^+(G)|, |\sigma^-(G)|\}$.

Dominating induced matchings and maximum induced matchings

- It is clear that if M is an induced matching of G , then

$$|M| \leq \frac{|E(G)|}{2\delta(G) - 1},$$

where $\delta(G)$ denotes the minimum degree of the vertices of G .

Dominating induced matchings and maximum induced matchings

- It is clear that if M is an induced matching of G , then

$$|M| \leq \frac{|E(G)|}{2\delta(G) - 1},$$

where $\delta(G)$ denotes the minimum degree of the vertices of G .

- If G is p -regular and M is a dominating induced matching, then the above upper bound is attained.

Dominating induced matchings and maximum induced matchings

- ▶ It is clear that if M is an induced matching of G , then

$$|M| \leq \frac{|E(G)|}{2\delta(G) - 1},$$

where $\delta(G)$ denotes the minimum degree of the vertices of G .

- ▶ If G is p -regular and M is a dominating induced matching, then the above upper bound is attained.
- ▶ Combining the above upper bound with the previous corollary, we may conclude that some graphs have no dominating induced matchings.

Graph eigenvalues and matchings

► Theorem[Ming and Wang, 2000]

Let T be a tree with n vertices. If T has a perfect matching then $\lambda_{n/2}(T) = 2$.

Graph eigenvalues and matchings

► Theorem[Ming and Wang, 2000]

Let T be a tree with n vertices. If T has a perfect matching then $\lambda_{n/2}(T) = 2$.

► Theorem[Brouwer and Haemers, 2005]

Let G be a connected k -regular graph with $2p$ vertices and with eigenvalues $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \dots \geq \lambda_{2p}$. If

$$\lambda_3 \leq \begin{cases} k - 1 + \frac{3}{k+1}, & \text{if } k \text{ is even} \\ k - 1 + \frac{3}{k+2}, & \text{otherwise,} \end{cases}$$

then G has a perfect matching.

Hamiltonian paths and Hamiltonian cycles

Considering a graph G of order n , in [Fiedler and Nikiforov, 2010] the following results are deduced.

- ▶ If $\lambda_1(G) \geq n - 2$ then G has a Hamiltonian path, unless $G = K_{n-1} + v$ (a complete graph plus an isolated vertex v).

Hamiltonian paths and Hamiltonian cycles

Considering a graph G of order n , in [Fiedler and Nikiforov, 2010] the following results are deduced.

- ▶ If $\lambda_1(G) \geq n - 2$ then G has a Hamiltonian path, unless $G = K_{n-1} + v$ (a complete graph plus an isolated vertex v).
- ▶ If $\lambda_1(G) > n - 2$ then G has a Hamiltonian cycle, unless $G = K_{n-1} + e$ (a complete graph plus a pendent edge e).

Hamiltonian paths and Hamiltonian cycles

Considering a graph G of order n , in [Fiedler and Nikiforov, 2010] the following results are deduced.

- ▶ If $\lambda_1(G) \geq n - 2$ then G has a Hamiltonian path, unless $G = K_{n-1} + v$ (a complete graph plus an isolated vertex v).
- ▶ If $\lambda_1(G) > n - 2$ then G has a Hamiltonian cycle, unless $G = K_{n-1} + e$ (a complete graph plus a pendent edge e).
- ▶ If $\lambda_1(\overline{G}) \leq \sqrt{n-1}$ then G has a Hamiltonian path, unless $G = K_{n-1} + v$.

Hamiltonian paths and Hamiltonian cycles

Considering a graph G of order n , in [Fiedler and Nikiforov, 2010] the following results are deduced.

- ▶ If $\lambda_1(G) \geq n - 2$ then G has a Hamiltonian path, unless $G = K_{n-1} + v$ (a complete graph plus an isolated vertex v).
- ▶ If $\lambda_1(G) > n - 2$ then G has a Hamiltonian cycle, unless $G = K_{n-1} + e$ (a complete graph plus a pendent edge e).
- ▶ If $\lambda_1(\overline{G}) \leq \sqrt{n-1}$ then G has a Hamiltonian path, unless $G = K_{n-1} + v$.
- ▶ If $\lambda_1(\overline{G}) \leq \sqrt{n-2}$ then G has a Hamiltonian cycle, unless $G = K_{n-1} + e$.

Definition of (k, τ) -regular set


A vertex subset $S \subseteq V(G)$ is (k, τ) -**regular** if induces a k -regular subgraph and

$$\forall v \notin S \quad |N_G(v) \cap S| = \tau.$$

Definition of (k, τ) -regular set

A vertex subset $S \subseteq V(G)$ is (k, τ) -**regular** if induces a k -regular subgraph and


$$\forall v \notin S \quad |N_G(v) \cap S| = \tau.$$



Definition of (k, τ) -regular set

A vertex subset $S \subseteq V(G)$ is (k, τ) -**regular** if induces a k -regular subgraph and

$$\forall v \notin S \quad |N_G(v) \cap S| = \tau.$$




$S_1 = \{1, 2, 3, 4\}$ is $(0, 2)$ -regular.

Definition of (k, τ) -regular set

A vertex subset $S \subseteq V(G)$ is (k, τ) -**regular** if induces a k -regular subgraph and

$$\forall v \notin S \quad |N_G(v) \cap S| = \tau.$$



$S_1 = \{1, 2, 3, 4\}$ is $(0, 2)$ -regular.

$S_2 = \{5, 6, 7, 8, 9, 10\}$ is $(1, 3)$ -regular.

Particular cases

- If G has a (k, τ) -regular set S then in its complement, \bar{G} , S is $(|S| - k - 1, |S| - \tau)$ -regular.

Particular cases

- ▶ If G has a (k, τ) -regular set S then in its complement, \bar{G}, S is $(|S| - k - 1, |S| - \tau)$ -regular.
- ▶ According to the definition, if a graph G is k -regular, then $V(G)$ is (k, τ) -regular for every nonnegative integer τ .

Particular cases

- ▶ If G has a (k, τ) -regular set S then in its complement, \bar{G}, S is $(|S| - k - 1, |S| - \tau)$ -regular.
- ▶ According to the definition, if a graph G is k -regular, then $V(G)$ is (k, τ) -regular for every nonnegative integer τ .
- ▶ By convention, if G is k -regular, then we say that $V(G)$ is $(k, 0)$ -regular.

Particular cases

- ▶ If G has a (k, τ) -regular set S then in its complement, \bar{G} , S is $(|S| - k - 1, |S| - \tau)$ -regular.
- ▶ According to the definition, if a graph G is k -regular, then $V(G)$ is (k, τ) -regular for every nonnegative integer τ .
- ▶ By convention, if G is k -regular, then we say that $V(G)$ is $(k, 0)$ -regular.
- ▶ If a p -regular graph G has a (k, τ) -regular set S , then $V(G) \setminus S$ is $(p - \tau, p - k)$ -regular.

Independent sets and dominating induced matchings

Theorem[Barbosa and C, 2004]

A graph G has a maximum stable set which is $(0, 1)$ -regular if and only if each vertex belongs to exactly one simplex.

Independent sets and dominating induced matchings

Theorem[Barbosa and C, 2004]

A graph G has a maximum stable set which is $(0, 1)$ -regular if and only if each vertex belongs to exactly one simplex.

Theorem[C, Cerdeira, Delorme and Silva, 2008]

If $M \subset E(G)$ is a dominated induced matching of G , then M is a maximum induced matching.

Independent sets and dominating induced matchings

Theorem[Barbosa and C, 2004]

A graph G has a maximum stable set which is $(0, 1)$ -regular if and only if each vertex belongs to exactly one simplex.

Theorem[C, Cerdeira, Delorme and Silva, 2008]

If $M \subset E(G)$ is a dominated induced matching of G , then M is a maximum induced matching.

A connected graph G with more than one edge has a dominating induced matching $D \subset E(G)$ if and only if $L(D)$ is $(0, 1)$ -regular in the line graph $L(G)$.

Independent sets

Theorem

Let G be a graph and $S \subset V(G)$. If S is $(0, \tau)$ -regular with $\tau = -\lambda_n(G)$, then S is a maximum stable set and every maximum stable set is $(0, \tau)$ -regular.

Independent sets

Theorem

Let G be a graph and $S \subset V(G)$. If S is $(0, \tau)$ -regular with $\tau = -\lambda_n(G)$, then S is a maximum stable set and every maximum stable set is $(0, \tau)$ -regular.



- ▶ The Petersen graph P has a $(0, 2)$ -regular set with cardinality 4 and $\lambda_n(P) = -2$. Then $\alpha(P) = 4$.
- ▶ The Hoffman-Singleton graph HS has a $(0, 3)$ -regular set with cardinality 15 and $\lambda_n(HS) = -3$. Then $\alpha(HS) = 15$.

Characterization of perfect matching

A connected graph G , with more than one edge, has a perfect matching M if and only if $L(M)$ is a $(0, 2)$ -regular set of the line graph $L(G)$.

Characterization of perfect matching

A connected graph G , with more than one edge, has a perfect matching M if and only if $L(M)$ is a $(0, 2)$ -regular set of the line graph $L(G)$.



Strongly regular graphs


Theorem [C, Sciriha and Zerafa, 2009]

A k -regular graph G of order n is strongly regular with parameters (n, k, a, c) if and only if $\forall v \in V(G)$, $S = N_G(v)$ is (a, c) -regular in $H = G - \{v\}$.


Strongly regular graphs

Theorem [C, Sciriha and Zerafa, 2009]

A k -regular graph G of order n is strongly regular with parameters (n, k, a, c) if and only if $\forall v \in V(G)$, $S = N_G(v)$ is (a, c) -regular in $H = G - \{v\}$.



The Petersen graph P



The graph $P - \{v\}$

Hamiltonian cycles


Lemma

A graph G has a Hamiltonian cycle C if and only if $L(C)$ is $(2, 4)$ -regular inducing a connected subgraph in the line graph $L(G)$.

Hamiltonian cycles

Lemma


A graph G has a Hamiltonian cycle C if and only if $L(C)$ is $(2, 4)$ -regular inducing a connected subgraph in the line graph $L(G)$.



Hamiltonian cycles

Lemma

A graph G has a Hamiltonian cycle C if and only if $L(C)$ is $(2, 4)$ -regular inducing a connected subgraph in the line graph $L(G)$.



Eigenvalues and (k, τ) -regular sets

Theorem[Thompson, 1981]

Let G be a p -regular graph and \mathbf{x} the characteristic vector of $S \subseteq V(G)$. Then S is a (k, τ) -regular set, with $\tau > 0$, if and only if $k - \tau \in \sigma(G)$ with corresponding eigenvector $(p - k + \tau)\mathbf{x} - \tau\mathbf{j}$, where \mathbf{j} is the all-one vector.

Eigenvalues and (k, τ) -regular sets


Theorem[Thompson, 1981]

Let G be a p -regular graph and \mathbf{x} the characteristic vector of $S \subseteq V(G)$. Then S is a (k, τ) -regular set, with $\tau > 0$, if and only if $k - \tau \in \sigma(G)$ with corresponding eigenvector $(p - k + \tau)\mathbf{x} - \tau \mathbf{j}$, where \mathbf{j} is the all-one vector.


Theorem[C and Rama, 2007]

Let $\lambda \in \mathbb{Z}$ and G be a graph with a (k_1, τ_1) -regular set S_1 ($\tau_1 > 0$) and a (k_2, τ_2) -regular set S_2 , such that $S_1 \neq S_2$ and $k_1 - \tau_1 = k_2 - \tau_2 = \lambda$. Then $\lambda \in \sigma(G)$ with corresponding eigenvector $\frac{\tau_2}{\tau_1} \mathbf{x}_1 - \mathbf{x}_2$, where \mathbf{x}_1 and \mathbf{x}_2 are the characteristic vectors of S_1 and S_2 , respectively.

Eigenvalues and eigenvectors




Eigenvalues and eigenvectors




$S_1 = \{1, 2, 5, 6\}$ and $S_2 = \{2, 3, 4, 5\}$ are $(2, 1)$ -regular.

Eigenvalues and eigenvectors



$S_1 = \{1, 2, 5, 6\}$ and $S_2 = \{2, 3, 4, 5\}$ are $(2, 1)$ -regular.
 $T_1 = \{1, 4\}$ and $T_2 = \{3, 6\}$ are $(0, 1)$ -regular.

Eigenvalues and eigenvectors



$S_1 = \{1, 2, 5, 6\}$ and $S_2 = \{2, 3, 4, 5\}$ are $(2, 1)$ -regular.

$T_1 = \{1, 4\}$ and $T_2 = \{3, 6\}$ are $(0, 1)$ -regular.

Then $\{-1, 1\} \subset \sigma(G) = \{-2.41, -1, -0.41, 0.41, 1, 2.41\}$ and

$$\begin{bmatrix} 1 \\ 0 \\ -1 \\ -1 \\ 0 \\ 1 \end{bmatrix} \in \mathcal{E}_G(1) \quad \text{and} \quad \begin{bmatrix} 1 \\ 0 \\ -1 \\ 1 \\ 0 \\ -1 \end{bmatrix} \in \mathcal{E}_G(-1).$$

Maximum induced k -regular subgraphs


Theorem[C, Kamiński and Lozin, 2007]

Let G be a graph and $\tau = -\lambda_n(G)$. If $S \subseteq V(G)$ is $(k, k + \tau)$ -regular, then S is a maximum cardinality set inducing a k -regular subgraph.

Maximum induced k -regular subgraphs

Theorem[C, Kamiński and Lozin, 2007]

Let G be a graph and $\tau = -\lambda_n(G)$. If $S \subseteq V(G)$ is $(k, k + \tau)$ -regular, then S is a maximum cardinality set inducing a k -regular subgraph.



Graph G with $\sigma(G) = \{-2, 0, 4\}$ and the $(2, 4)$ -regular set $\{1, 3, 4, 6\}$.

The main characteristic polynomial

Definition

Each distinct eigenvalue μ_1, \dots, μ_p , $1 \leq p \leq n$, of a graph G such that $\mathcal{E}_G(\mu_i)$ is *not* orthogonal to the all-one vector \mathbf{j} is said to be **main**. The remaining distinct eigenvalues are **non-main**.

The main characteristic polynomial

Definition

Each distinct eigenvalue μ_1, \dots, μ_p , $1 \leq p \leq n$, of a graph G such that $\mathcal{E}_G(\mu_i)$ is *not* orthogonal to the all-one vector \mathbf{j} is said to be **main**. The remaining distinct eigenvalues are **non-main**.

Lemma [D. Cvetković and M. Petrić, 1984]

If G is a graph with p distinct main eigenvalues μ_1, \dots, μ_p , then the **main characteristic polynomial** of G

$$m_G(\lambda) = \lambda^p - c_0 \lambda^{p-1} - c_1 \lambda^{p-2} - \dots - c_{p-2} \lambda - c_{p-1}$$

has integer coefficients.

The walk matrix


Definition

Given a graph G of order n , the $n \times k$ walk matrix of G is the matrix $\mathbf{W}' = (\mathbf{j}, \mathbf{A}_G \mathbf{j}, \mathbf{A}_G^2 \mathbf{j}, \dots, \mathbf{A}_G^{k-1} \mathbf{j})$.

The walk matrix

Definition

Given a graph G of order n , the $n \times k$ walk matrix of G is the matrix $\mathbf{W}' = (\mathbf{j}, \mathbf{A}_G \mathbf{j}, \mathbf{A}_G^2 \mathbf{j}, \dots, \mathbf{A}_G^{k-1} \mathbf{j})$.



$$\mathbf{W}' = (\mathbf{j}, \mathbf{A}_G \mathbf{j}, \mathbf{A}_G^2 \mathbf{j})$$

$$= \begin{pmatrix} 1 & 3 & 11 \\ 1 & 4 & 12 \\ 1 & 4 & 12 \\ 1 & 3 & 11 \\ 1 & 2 & 6 \\ 1 & 2 & 6 \end{pmatrix}$$

The column space of the walking matrix

Theorem (Hagos, 2002)

Let G be a graph of order n with p distinct main eigenvalues.
The rank of its $n \times k$ walk matrix W' , is p for $k \geq p$.

The column space of the walking matrix

Theorem (Hagos, 2002)

Let G be a graph of order n with p distinct main eigenvalues. The rank of its $n \times k$ walk matrix W' , is p for $k \geq p$.

Then the number of distinct main eigenvalues p is such that $p = \min\{k : \{j, Aj, A^2j, \dots, A^kj\} \text{ is linear dependent}\}$.

The column space of the walking matrix

Theorem (Hagos, 2002)

Let G be a graph of order n with p distinct main eigenvalues. The rank of its $n \times k$ walk matrix W' , is p for $k \geq p$.

Then the number of distinct main eigenvalues p is such that $p = \min\{k : \{j, Aj, A^2j, \dots, A^kj\} \text{ is linear dependent}\}$.

The p th column of AW is $A^p j = W \begin{pmatrix} c_{p-1} \\ \vdots \\ c_1 \\ c_0 \end{pmatrix}$, where c_j , for $j = 0, \dots, p-1$, are the coefficients of $m_G(\lambda)$.

Main (non-main) eigenvalues of graphs with (k, τ) -regular sets

Definition

The main eigenspace of a graph G , $Main(G)$, is the subspace spanned by the eigenvectors not orthogonal to the all one vector j .

Main (non-main) eigenvalues of graphs with (k, τ) -regular sets

Definition

The main eigenspace of a graph G , $\text{Main}(G)$, is the subspace spanned by the eigenvectors not orthogonal to the all one vector \mathbf{j} .

Theorem[C, Sciriha and Zerafa, 2008]

Let G be a graph with a (k, τ) -regular set S , where $\tau > 0$, and $\lambda \in \sigma(G)$.

1. The eigenvalue λ is non-main if and only if

$$\lambda = k - \tau \quad \text{or} \quad \mathbf{x}_S \in (\mathcal{E}_G(\lambda))^\perp.$$
2. If λ is main with associated eigenvector $\mathbf{u} \in \text{Main}(G)$, then

$$\mathbf{u}^t \mathbf{x}_S \neq 0 \text{ and } \lambda = \tau \frac{\mathbf{u}^t \mathbf{x}_S}{\mathbf{u}^t \mathbf{x}_S} + k.$$

(κ, τ) -parametric vectors

Definition

Let G be a graph with p distinct main eigenvalues μ_1, \dots, μ_p and $\{x_1, \dots, x_p\}$ an orthonormal basis of $\text{Main}(G)$. Considering the nonnegative integer κ and the positive integer τ such that $(\kappa - \tau) \notin \{\mu_1, \dots, \mu_p\}$, the vector

$$g = \sum_{i=1}^p \tau \frac{j^t x_i}{\mu_i - (\kappa - \tau)} x_i$$

is referred as the (κ, τ) -parametric vector of G .

Graphs with (κ, τ) -regular sets

Theorem[C, Sciriha and Zerafa, 2008]

Let G be a graph with p distinct main eigenvalues μ_1, \dots, μ_p and $\{x_1, \dots, x_p\}$ an orthonormal basis of $\text{Main}(G)$. Consider a (κ, τ) -regular set $S \subseteq V(G)$, with $\tau > 0$. Then the characteristic vector of S , x_S , is such that

$$x_S = g + q,$$

where $q \in (\text{Main}(G))^\perp$ and the following holds.

1. If $\kappa - \tau \notin \sigma(G)$, then $q = 0$ and $x_S = \tau (A - (\kappa - \tau)I)^{-1} j$;
2. If $\kappa - \tau \in \sigma(G)$ then $q \in \varepsilon_G(\kappa - \tau)$.

Graphs with (κ, τ) -regular sets

Corollary[C, Sciriha and Zerafa, 2008]

If a graph G with p distinct main eigenvalues μ_1, \dots, μ_p and $\{x_1, \dots, x_p\}$ an orthonormal basis of $\text{Main}(G)$ has a (κ, τ) -regular set $S \subset V(G)$ and $(\kappa - \tau) \in \sigma(G)$, then

$$|S| = \sum_{i=1}^p \tau \frac{(j^t x_i)^2}{\mu_i - (\kappa - \tau)}.$$

Bibliography



R. BARBOSA AND D. M. CARDOSO, "On regular-stable graphs," *ARS Combinatoria* 70 (2004), 149-159.



D. M. Cardoso, P. Rama, "Equitable bipartitions of graphs and related results," *J. Math. Sciences* **120** (Issue 1: Aveiro Seminar on Control, Optimization and Graph Theory) (2004), 869-880.



D. M. CARDOSO AND D. M. CVETKOVIĆ, "Graphs with least eigenvalue -2 attaining a convex quadratic upper bound for the stability number," *Bull. Acad. Serbe Sci. Arts - 2006, Cl. Sci. Math. Natur., Sci. Math.*, 81 (2006): 41-55.



D. M. CARDOSO, M. KAMINSKI AND V. LOZIN, "Maximum k -regular induced subgraphs," *J. Combin. Optim.*, Vol. 14, n. 4 (2007): 455-463.

Bibliography



D. M. CARDOSO AND P. RAMA, "Spectral results on graphs with regularity constraints," *Linear Algebra Appl.* **423** (2007), 90-98.



D. M. CARDOSO, J. O. CERDEIRA, C. DELORME, P. C. SILVA, "Efficient edge domination in regular graphs," *Discrete Applied Math.* **156** (2008), 3060-3065.



D. M. CARDOSO, I. SCIRIHA AND C. ZERAFA, "Main eigenvalues and (k, τ) -regular sets", *Linear Algebra Appl.* **432** (2010):2399-2408.



D. M. CARDOSO AND P. ROWLINSON, "Spectral upper bounds for the order of a k -regular induced subgraph", *Linear Algebra Appl.* **433** (2010):1031-1037.

Bibliography



D. CVETKOVIĆ, "Inequalities obtained on the basis of the spectrum of the graph," *Studia Sci. Math. Hung.* 8 (1973): 433-436.



D. CVETKOVIĆ, M. DOOB AND H. SACHS, "Spectra of Graphs," Academic Press, New York, 1979.



D. CVETKOVIĆ, "Chromatic number and the spectrum of a graph," *Publ. Inst. Math. (Beograd)* 14 (28) (1972): 25-38.



D. CVETKOVIĆ AND M. PETRIĆ, "A Table of Connected Graphs on Six Vertices," *Discrete Math.* **50**-1 (1984): 37-49.



M. FIEDLER AND V. NIKIFOROV, "Spectral radius and Hamiltonicity of graphs," *Linear Algebra Appl.* **432** (2010): 2170-2173.

Bibliography



G. FINKE, R. E. BURKARD AND F. RENDL, *Quadratic assignment problem*, Surveys in combinatorial optimization (Rio de Janeiro, 1985): 61–82, North-Holland Math. Stud., 132, North-holland, Amsterdam, 1987.



E.M., HAGOS, "Some results on graph spectra," *Linear Algebra Appl.* **356** (2002): 103-111.



M. M. HALLDÓRSSON, J. KRATOCHVÍL AND J. A. TELLE, "Independent sets with domination constraints," *Discrete Applied Mathematics*, 99 (2000): 39–54.



A. HOFFMAN, "On eigenvalues and colorings of graphs," in "Graph Theory and its Applications" (B. Harris, ed.), Academic Press, New York (1970): 79-91.

Bibliography



Y. HOU AND F. TIAN, "Unicyclic graphs with exactly two main eigenvalues," *Applied Mathematics Letters* **19** (2005): 1143-1147.



T. MOTZKIN AND E. G. STRAUS, "Maxima for graphs and new proof of a theorem of trán," *Can. J. Math.* **17** (1965): 533-540.



P., ROWLINSON, "The Main Eigenvalues of a Graph: A Survey," *Applicable Analysis and Discrete Mathematics*, **1** (2007): 445-471.



A. NEUMAIER, "Regular sets and quasi-symmetric 2-designs," in: D. Jungnickel and K. Vedder (Eds.), *Combinatorial Theory*, Lecture Notes in Math., 969, Springer-Verlag, Berlin, (1982): 258–275.

Bibliography



R. RYACEK AND I. SCIERMEYER, "On the independence number of $K_{1,r+1}$ -free graphs," *Discrete Mathematics* 138 (1995): 265-374.



J. A. TELLE, "Characterization of domination-type parameters in graphs, *Congressus Numerantium*, 94 (1993): 9–16.



D. M. THOMPSON, "Eigengraphs: constructing strongly regular graphs with block designs," *Utilitas Mathematica*, 20 (1981): 83–115.



H. S. WILF, "The eigenvalues of a graph and its chromatic number," *J. London Math. Soc.* 42 (1967): 330-332.