



Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Constructing infinite families of ALQ -integral graphs

Nair Abreu

Production Engineering Program

Federal University of Rio de Janeiro, Brazil

ILAS, Pisa, Italy - June, 2010.



Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

My thanks to Vlado Nikiforov and Dragan Stevanović for their kind invitation to talk here!

My talk is based on a joint paper with Maria Agueiras A. de Freitas, Renata Del Vecchio and Cybele Maia Vinagre. My thanks to my coauthors!

Many thanks to the Brazilian Council for Scientific and Technological Development for support received for our research.



Topics

Basic
concepts and
results

Basic concepts and results;

A new
operation and
 ALQ -integral
graphs

A new operation and ALQ -integral graphs;

ALQ -integral
graphs and
regular graphs

ALQ -integral graphs and regular graphs;

ALQ -integral
graphs and
split-like
graphs

ALQ -integral graphs and split-like graphs.



Basic definitions

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

- $G = (V, E)$ is a simple and connected graph on n vertices and m edges;
- $A(G)$ is the adjacency matrix of G and $\lambda_1 \geq \dots \geq \lambda_n$ its eigenvalues;
- $D(G)$ is the diagonal matrix where the elements on the main diagonal are the degrees of the vertices of G ;
- $L(G) = D(G) - A(G)$ is the Laplacian matrix of G and $\mu_1 \geq \dots \geq \mu_{n-1} \geq \mu_n = 0$ its eigenvalues;
- $Q(G) = D(G) + A(G)$ is the signless Laplacian matrix of G and $q_1 \geq \dots \geq q_n$ its eigenvalues.



M -integral graphs

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Definition: Let M be a matrix associated to a graph G . A graph is M -integral if and only if the eigenvalues of M are all integer numbers.

- If $M = A$ is the adjacency matrix of G , G is called an A -integral graph;
- If $M = L$ is the Laplacian matrix of G , G is called an L -integral graph;
- If $M = Q$ is the Signless Laplacian matrix of G , G is called a Q -integral graph.



Regular graphs and integrality

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Theorem: If G is an r -regular graph, then

$$P_A(G, x) = P_Q(G, x + r) \text{ and } P_L(G, x) = (-1)^n P_Q(G, 2r - x).$$

Consequently,

$$P_L(G, x) = (-1)^n P_A(G, r - x).$$

Corollary: Let G be a regular graph. G is an A -integral graph
 $\Leftrightarrow G$ is an L -integral graph $\Leftrightarrow G$ is a Q -integral graph.



Bipartite graphs and integrality

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Theorem: Let G be a bipartite graph. G is an L -integral graph $\Leftrightarrow G$ is a Q -integral graph.



ALQ -integral graphs

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Definition: *A graph is called an ALQ -integral when it is simultaneously A , L and Q -integral graph.*

- In 2007, Z. Stanić enumerated all connected Q -integral graphs up to 10 vertices.
- There are 172 such graphs.
- Among them, there are 42 ALQ -integral graphs.
- It is interesting to note that only one of them is neither regular nor bipartite.



The smallest connected neither bipartite nor regular ALQ -integral graph

There are no connected neither regular nor bipartite ALQ -integral graph up to 9 vertices.




Figure: The smallest connected ALQ -integral graph (it has 10 vertices) which is neither bipartite nor regular.

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs



Objective

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

- **Our aim:** To construct infinite families of ALQ -integral graphs which necessarily contain neither regular nor bipartite graphs.
- **Remark:** If we consider the computational experiments of Stanić(2007), we can hope that the ALQ -integral graphs, neither bipartite nor regular, have huge orders. So, we are looking for them through the operation of graphs.



Union and Cartesian product of graphs

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Let G_1 and G_2 be graphs such that, for $i = 1, 2$, V_i and E_i are, respectively, the sets of vertices and edges of G_i .

Union of graphs: *The union $G_1 \cup G_2$ of G_1 and G_2 is the graph whose vertex set is $V = V_1 \cup V_2$ and whose edge set is $E = E_1 \cup E_2$.*

Cartesian Product: *The Cartesian Product $G_1 \times G_2$ of G_1 and G_2 is the graph whose vertex set is $V = V_1 \times V_2$ and where*

$$(u_1, u_2) \sim (v_1, v_2) \Leftrightarrow \left\{ \begin{array}{l} u_1 \sim v_1 \text{ in } G_1 \text{ and } u_2 = v_2, \text{ or} \\ u_1 = v_1 \text{ and } u_2 \sim v_2 \text{ in } G_2. \end{array} \right\}$$



Spectra of resultant graphs from operations

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

- G_1 and G_2 are A -integral graphs $\Rightarrow G_1 \cup G_2$ and $G_1 \times G_2$ are A -integral graphs;
- G_1 and G_2 are L -integral graphs $\Rightarrow G_1 \cup G_2$ and $G_1 \times G_2$ are L -integral graphs;
- G_1 and G_2 are Q -integral graphs $\Rightarrow G_1 \cup G_2$ and $G_1 \times G_2$ are Q -integral graphs.



A new operation

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Definition of k -pla operation: Let $G = (V, E)$ be a graph on n vertices and, for $k \in \mathbb{N}$, let $K = \{1, \dots, k\} \subset \mathbb{N}$. We call a k -pla graph $G^{(k)} = (V_k, E_k)$ of G that one which set of vertices is $V_k = V \times K$ and the set of edges is $E_k = \{\{(x_1, y_1), (x_2, y_2)\}; \{x_1, x_2\} \in E; y_1, y_2 \in K\}$.

Remark:

- This definition generalizes the definition of a double graph of G , given by Munarini *et al* (2008). In fact, the double graph is a 2-pla graph of G .
- It is a special case of NEPS (see Cvetković *et al.*)



An example of a 3-pla graph

Basic
concepts and
results

**A new
operation and
 ALQ -integral
graphs**

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs




Figure: 3-pla graph of P_3



Matrices of a k -pla graph

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

- Note that $G^{(1)} \simeq G$.
- The adjacency matrix of $G^{(k)}$ can be represented as

$$A(G^{(k)}) = \mathbb{J}_k \otimes A(G),$$

where \mathbb{J}_k denotes the all ones matrix of order k and \otimes is the Kronecker product of matrices.

- Consequently,

$$L(G^{(k)}) = k \mathbb{I}_k \otimes D(G) - \mathbb{J}_k \otimes A(G)$$

and

$$Q(G^{(k)}) = k \mathbb{I}_k \otimes D(G) + \mathbb{J}_k \otimes A(G).$$



The spectra of a k -pla graph

Theorem: Let G be a graph on n vertices and $D(G) = \text{diag}(d_1, \dots, d_n)$, $d_i, 1 \leq i, n$, is degree of the vertex v_i in G . Let $k \in \mathbb{N}$, $k \geq 2$.

- The A -eigenvalues of $G^{(k)}$ are $0^{(k-1)n}, k\lambda_1, \dots, k\lambda_n$, where $Sp_A(G) = (\lambda_1, \dots, \lambda_n)$;
- The L -eigenvalues of $G^{(k)}$ are $nd_1^{k-1}, \dots, nd_n^{k-1}, k\mu_1, \dots, k\mu_n$, where $Sp_L(G) = (\mu_1, \dots, \mu_n)$;
- The Q -eigenvalues of $G^{(k)}$ are $nd_1^{k-1}, \dots, kd_n^{k-1}, kq_1, \dots, kq_n$, where $Sp_Q(G) = (q_1, \dots, q_n)$.

Observation: The exponents denote multiplicities.

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs



Sketch of the Proof

Sketch of the Proof: We have,

- $A(G)\mathbf{v} = \lambda\mathbf{v} \implies A(G^{(k)})\mathbf{w} = (\mathbb{J}_k \otimes A(G))\mathbf{1}_k \otimes \mathbf{v} = k\lambda\mathbf{w}$, where $\mathbf{w} = \mathbf{1}_k \otimes \mathbf{v}$.
- $\mathbf{u} \in \mathbb{R}^k$ is orthogonal to $\mathbf{1}_k \implies A(G^{(k)})\mathbf{w}_i = \mathbf{0}$, where $\mathbf{w}_i = \mathbf{u} \otimes \mathbf{e}_i$, for $1 \leq i \leq n$.

Consequently, the A -eigenvalues of $G^{(k)}$ are

$$0^{(k-1)n}, k\lambda_1, \dots, k\lambda_n,$$

where $Sp_A(G) = (\lambda_1, \dots, \lambda_n)$. ■

Observation: The proofs of the others items are similar.



ALQ -integrality condition for a k -pla graph

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Corollary: *The graph $G^{(k)}$, $k \in \mathbb{N}$, $k \geq 2$, is ALQ -integral if and only if G is an ALQ -integral graph.*



Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Corollary: *For each ALQ -integral graph G , $\{G^{(k)}, k \in \mathbb{N}\}$ is an infinite family of ALQ -integral graphs.*

Remark: Note that if G is neither regular nor bipartite, the same occurs with $G^{(k)}$, for each $k \in \mathbb{N}$.



Join of graphs

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Definition of Join Operation: *Let G_1 and G_2 be graphs such that, for $i = 1, 2$, V_i and E_i are, respectively, the sets of vertices and edges of G_i . The Join $G_1 \vee G_2$ of G_1 and G_2 is the graph obtained from $G_1 \cup G_2$ by joining each vertex of G_1 with every vertex of G_2 .*



The L -characteristic polynomial of a join of graphs

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Theorem: For $i = 1, 2$, let G_i be a graph on n_i vertices.
Then, the L -characteristic polynomial of $G_1 \vee G_2$ is

$$P_L(G_1 \vee G_2, x) = x(x - (n_1 + n_2)) \frac{P_L(G_1, x - n_2)P_L(G_2, x - n_1)}{(x - n_2)(x - n_1)}. \quad (1)$$

Corollary: $G_1 \vee G_2$ is L -integral $\Leftrightarrow G_1$ and G_2 are L -integral.



The A -characteristic polynomial of a join of regular graphs

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Theorem: For $i = 1, 2$, let G_i be a r_i -regular graph on n_i vertices. Then, the A -characteristic polynomial of $G_1 \vee G_2$ is

$$P_A(G_1 \vee G_2, x) = \frac{P_A(G_1, x)P_A(G_2, x)}{(x - r_1)(x - r_2)} f(x), \quad (2)$$

where $f(x) = x^2 - (r_1 + r_2)x + r_1r_2 - n_1n_2$.



The Q -polynomial of a join of regular graphs

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Theorem(Freitas et al., 2010) For $i = 1, 2$, let G_i be a r_i -regular graph on n_i vertices. Then, the Q -characteristic polynomial of $G_1 \vee G_2$ is

$$P_Q(G_1 \vee G_2, x) = \frac{P_Q(G_1, x - n_2)P_Q(G_2, x - n_1)}{(x - 2r_1 - n_2)(x - 2r_2 - n_1)} f(x), \quad (3)$$

where

$$f(x) = x^2 - (2(r_1 + r_2) + (n_1 + n_2))x + 2(2r_1r_2 + r_1n_1 + r_2n_2).$$



ALQ -integrality condition for a join of regular graphs

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Theorem: For $i = 1, 2$, let G_i be a r_i -regular graph on n_i vertices. The graph $G_1 \vee G_2$ is ALQ -integral if and only if G_1 and G_2 are ALQ -integral and $(r_1 - r_2)^2 + 4n_1n_2$ and $((2r_1 - n_1) - (2r_2 - n_2))^2 + 4n_1n_2$ are perfect squares.

Note that if $r_1 - r_2 \neq 0$ and $r_1 - r_2 \neq n_1 - n_2$ then $G_1 \vee G_2$ is neither regular nor bipartite.

From this result, we obtain some corollaries which give us more infinite families of ALQ -integral graphs.



$K_2 \vee 4K_2$ satisfies the last theorem


The smallest connected neither bipartite nor regular ALQ -integral graph is isomorphic to $K_2 \vee 4K_2$ and it satisfies the conditions of the last theorem.

$G_1 = K_2$ has 2 vertices and $G_2 = 4K_2$ has 8 vertices and both graphs are 1-regular;

$$(r_1 - r_2)^2 + 4n_1n_2 = 64 \text{ is a perfect square;}$$

$$((2r_1 - n_1) - (2r_2 - n_2))^2 + 4n_1n_2 = 36;$$

Also, we have $r_1 - r_2 = 0 \neq n_2 - n_1 = 6$.





The first corollary

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Corollary I: *Let G_1 and G_2 be r -regular ALQ -integral graphs. Then, $G_1 \vee G_2$ is ALQ -integral if and only if $|G_1||G_2|$ is a perfect square.*

Sketch of the Proof: Let $|G_i| = n_i$, $i = 1, 2$, both regular graphs with the same degree r . From last proposition, we have:

$G_1 \vee G_2$ is ALQ -integral $\Leftrightarrow ((2r - n_1) - (2r - n_2))^2 + 4n_1n_2 = (n_1 + n_2)^2$ and $4n_1n_2$ are perfect squares.

So, $G_1 \vee G_2$ is ALQ -integral $\Leftrightarrow |G_1||G_2|$ is a perfect square.





The first corollary generalizes a result of Stanić(2007)

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Lemma: A complete bipartite graph $K_{t,s}$ is ALQ -integral if and only if ts is a perfect square.



The second corollary

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Corollary II: *For all $j, k \in \mathbb{N}$, the graph $K_j \vee kK_j$ is ALQ -integral if and only if k is a perfect square.*

The second corollary also generalizes the other result of Stanić for $j = 2$. His result was recently published in *Ars Combinatoria* (2009).



The third corollary

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Corollary III: *For all $j \in \mathbb{N}$ and $k = \frac{j(j+1)}{2}$, the graph $K_{j,j} \vee kK_{j+1}$ is ALQ -integral.*



An example which satisfies the third corollary

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs




Figure: $K_{2,2} \vee 3K_3$



Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

In 2002, Hansen *et al* characterized A -integral graphs and, in 2010, Freitas *et al* characterized Q -integral graphs in the classes constituted by joins of complete graphs with the complements of complete graphs.

Let $a, b, k \in \mathbb{N}$. They are the following classes:

Complete split graphs, $CS_b^a \cong \overline{K_a} \vee K_b$;

Multiple complete split-like graphs,

$MCS_{b,k}^a \cong \overline{K_a} \vee (kK_b)$;

Multiple extended complete split-like graphs,

$MECS_{b,k}^a \cong \overline{K_a} \vee (k(K_b \times K_2))$.



ALQ -integrality condition for split like graphs

Corollary 4: For $a, b \in \mathbb{N}$ and $k \in \mathbb{N}$,

the complete split graph CS_b^a is ALQ -integral if and only if $(b-1)^2 + 4ab$ and $(a+b-2)^2 + 4ab$ are perfect squares;

the multiple complete split-like graph $MCS_{b,k}^a$ is ALQ -integral if and only if $(a+2(b-1)-kb)^2 + 4abk$ and $(b-1)^2 + 4abk$ are perfect squares;

the multiple extended complete split-like graph $MECS_{b,k}^a$ is ALQ -integral if and only if $(a+2b(k-1))^2 + 8ab$ and $b^2 + 8abk$ are perfect squares.

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs



Sketch of the Proof:

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Sketch of the Proof of the first item: $\overline{K_a}$ and K_b are regular ALQ -integral graphs of orders a and b and degrees 0 and $b - 1$, respectively.

From the last corollary, CS_b^a is ALQ -integral $\Leftrightarrow (b - 1)^2 + 4ab$ and $(a + b - 2)^2 + 4ab$ are perfect squares. ■

Observation: The proof of the others items are similar.



Families of ALQ -integral complete split graphs

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Can we determine some graphs which satisfy the conditions of the last theorem?



Some examples: If one of the conditions below holds, CS_b^a is Q -integral:

$a = 3j_t$ and $b = 2j_t - 1$, where $j_0 = 3$, $m_0 = 7$ and there is $m_t \in \mathbb{Z}$, $t \geq 1$ such that

$$\begin{aligned}j_{t+1} &= 127j_t + 484m_t - 45 \\m_{t+1} &= 336j_t + 127m_t - 120.\end{aligned}$$

$a = 10j_t - 4$, $b = 3j_t$, where $j_0 = 4$, $m_0 = 43$ and there is $m_t \in \mathbb{Z}$ such that

$$\begin{aligned}j_{t+1} &= -16855j_t - 1484m_t + 3528 \\m_{t+1} &= -191436j_t - 16855m_t + 40068;\end{aligned}$$

$a = 3j_t - 2$ and $b = 2j_t$, where $j_0 = 10$, $m_0 = -51$ and there is $m_t \in \mathbb{Z}$ such that

$$\begin{aligned}j_{t+1} &= 127j_t + 24m_t - 45 \\m_{t+1} &= 672j_t + 127m_t - 240;\end{aligned}$$

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs



Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Observation: As we built the families of ALQ -integral graphs for the complete split graphs, we can construct the infinite families for the multiple complete split graphs and the multiple extended complete split graphs in the similar way.

A complete split graph and its spectra

Figure 1 shows the complete split graph CS_5^9 whose spectra relative to A , L and Q respectively are

$$\begin{aligned}Sp_A(CS_5^9) &= (9, 0^8, -1^4, -5), \\Sp_L(CS_5^9) &= (14^5, 5^8, 0) \text{ and} \\Sp_Q(CS_5^9) &= (20, 12^4, 5^8, 2).\end{aligned}$$




Figure: CS_5^9

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs



A multiple complete split graph and its spectra

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs



Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

Figure below displays the multiple complete split graph, $MCS_{3,3}^7$, whose spectra relative to A , L and Q respectively are:

$$Sp_A(MCS_{3,3}^7) = (9, 2^2, 0^6, -1^6, -7),$$

$$Sp_L(MCS_{3,3}^7) = (16, 10^6, 9^6, 7^2, 0) \text{ and}$$

$$Sp_Q(MCS_{3,3}^7) = (18, 11^2, 9^6, 8^6, 2).$$




Figure: The graph $MCS_{3,3}^7$ is ALQ -integral

A multiple extended complete split graph and its spectra

Example: Figure below shows $G = MECS_{2,2}^6$ whose spectra

relative to A , L and Q are:

$$Sp_A(G) = (8, 2, 0^9, -2^2, -6),$$

$$Sp_L(G) = (14, 10^2, 8^9, 6, 0)$$

$$Sp_Q(G) = (16, 10, 8^9, 6^2, 2).$$




Figure: The graph $MECS_{2,2}^6$ is ALQ -integral

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs











Bibliography

Basic
concepts and
results

A new
operation and
 ALQ -integral
graphs

ALQ -integral
graphs and
regular graphs

ALQ -integral
graphs and
split-like
graphs

-  F. Harary, A.J. Schwenk, *Which graphs have integral spectra?*, em Bari, R., Harary, F.(Eds), Graphs and Combinatorics, Springer, Berlim, 1974, p. 45-51.
-  D. Cvetković, P. Rowlinson, S. Simić, *Signless Laplacian of finite graphs*, Linear Algebra and its Applications **423** (2007) 155-171.
-  S. Simić, Z. Stanić, *Q -integral graphs with edge-degrees at most five*, Discrete Math. **308** (2008) 4625-4634.
-  Z. Stanić, *There are exactly 172 connected Q -integral graphs up to 10 vertices*, Novi Sad J. Math. **37** (2) (2007) 193-205.
-  Stanić, Z., *Some results on Q -integral graphs*, Ars Combinatoria **90** (2009), 321-335.
-  M.A.A. de Freitas, N.M.M. de Abreu, R.R. Del-Vecchio, S.Jurkiewicz, *Infinite families of Q -integral graphs*, Linear Algebra Appl. **432** (2010), 2353-2360
-  D. Cvetković, M. Doob, H. Sachs: *Spectra of graphs-theory and application*. Deutscher Verlag der Wissenschaften-Academic Press, Berlin-New York, 1980; second edition 1982; third edition, Johann Ambrosius Barth Verlag, Heidelberg-Leipzig, 1995.
-  P. Hansen, H. Melot, D, Stevanović, *Integral complete split graphs*, Univ. Beograd, *Publ. Elektrotehn. Fak. Ser. Mat.* **13**, 2002, 89-95.